

Research Article

# Whole genome survey analysis and microsatellite motif identification of *Sebastiscus marmoratus*

 Sheng-yong Xu<sup>1</sup>, Na Song<sup>2</sup>, Shi-jun Xiao<sup>3</sup> and  Tian-xiang Gao<sup>1</sup>

<sup>1</sup>Fishery College, Zhejiang Ocean University, Zhoushan 316022, P.R. China; <sup>2</sup>Institute of Evolution and Marine Biodiversity, Ocean University of China, Qingdao 266003, P.R. China; <sup>3</sup>School of Computer Science and Technology, Wuhan University of Technology, Wuhan 430070, P.R. China

**Correspondence:** Tian-xiang Gao (gaotianxiang0611@163.com)



The marbled rockfish *Sebastiscus marmoratus* is an ecologically and economically important marine fish species distributed along the northwestern Pacific coast from Japan to the Philippines. Here, next-generation sequencing was used to generate a whole genome survey dataset to provide fundamental information of its genome and develop genome-wide microsatellite markers for *S. marmoratus*. The genome size of *S. marmoratus* was estimated as approximate 800 Mb by using K-mer analyses, and its heterozygosity ratio and repeat sequence ratio were 0.17% and 39.65%, respectively. The preliminary assembled genome was nearly 609 Mb with GC content of 41.3%, and the data were used to develop microsatellite markers. A total of 191,592 microsatellite motifs were identified. The most frequent repeat motif was dinucleotide with a frequency of 76.10%, followed by 19.63% trinucleotide, 3.91% tetranucleotide, and 0.36% pentanucleotide motifs. The AC, GAG, and ATAG repeats were the most abundant motifs of dinucleotide, trinucleotide, and tetranucleotide motifs, respectively. In summary, a wide range of candidate microsatellite markers were identified and characterized in the present study using genome survey analysis. High-quality whole genome sequence based on the “Illumina+PacBio+Hi-C” strategy is warranted for further comparative genomics and evolutionary biology studies in this species.

## Introduction

The assessment of genetic diversity and structure is one of the major goals of population management and conservation biology [1]. This assessment should ideally be achieved by utilizing polymorphic and informative markers. Microsatellites or simple sequence repeats (SSRs) are short tandem repeated motif (1–6 bases) that are found in both non-coding and coding regions of the genome and are characterized by a high degree of length polymorphism [2]. Microsatellite markers have become one of the most popular molecular markers and have been widely used in genetic studies due to their ubiquitous occurrence, high reproducibility, multiallelic nature, and codominant mode [2,3]. The advantages of microsatellite markers have made them one of the most useful tools for detecting genetic diversity, genetic linkage mapping, genetic structure, and germplasm and evolution analysis. However, conventional approaches to isolate and develop microsatellite primers were time- and cost-consuming because it is necessary to create enriched microsatellite libraries [2]. Until recently, next-generation sequencing (NGS) has provided a new perspective for the development of studies of microsatellite markers, owing to its high throughput and speed of data generation. So far, NGS has been applied to genomics-based strategies to discover sequences for new microsatellite markers in animals and plants, in a time- and cost-effective manner [4–8]. Genome survey sequencing (GSS) based on the NGS platform has been proven particularly useful in identifying genome-wide microsatellite markers in non-model species. Microsatellite markers development studies from GSS were performed in numbers of species [9–13]. Genome survey studies also provide information about genome structure of organisms, including estimates of genome size, levels of heterozygosity, and repeat contents.

Received: 18 July 2019  
Revised: 04 February 2020  
Accepted: 13 February 2020

Accepted Manuscript online:  
14 February 2020  
Version of Record published:  
24 February 2020

**Table 1** Quality control information of Illumina sequencing data

Lib ID	Raw data (bp)	Clean data (bp)	Effective rate (%)	Error rate (%)	Q20	Q30	GC content (%)
DES.L5	35,057,094,600	34,843,246,022	99.39	0.02; 0.03	98.02; 95.13	94.91; 89.18	42.86; 43.04

Note: The two statistics of error rate, Q20, Q30, and GC content were for pair-end read 1 and read 2, respectively.

The marbled rockfish (*Sebastes marmoratus*, Cuvier, 1829) is an ecologically and economically important ovoviparous marine species inhabiting littoral rocky bottoms along the northwest Pacific coast from Japan to the Philippines [14]. *S. marmoratus* has strong site fidelity and appears within narrow home ranges [14]. Several studies have been conducted on *S. marmoratus* germplasm resources due to the decline in wild populations [15–17]. However, inconsistent results were demonstrated given the insufficient resolution of molecular markers. Till now, limited microsatellite marker resources are publically available for *S. marmoratus* using different methods [18–22]. The use of microsatellite markers in molecular studies is limited and more microsatellite markers are needed for further studies. In the present study, we aimed to characterize and develop genome-wide microsatellite markers in *S. marmoratus* by genome survey sequencing. The newly identified microsatellites would be useful for extending our current knowledge of *S. marmoratus* genome organization and for genome mapping, marker-aided selection, and population genetics.

## Materials and methods

### Sample collection and genome survey sequencing

One male adult *S. marmoratus* was collected from Rushan (36°43'N, 121°39'E), China in October 2015. Muscle tissue was stored in 95% ethanol at –80°C. Total genomic DNA was extracted using a standard phenol–chloroform method for muscle tissue. DNA was treated with RNase A to produce pure, RNA-free DNA. Two paired-end DNA libraries were constructed with insert size of 350 bp, and then sequenced using the Illumina HiSeq2500 platform following the manufacturer's protocol. The library construction and sequencing were performed at Novogene in Beijing.

### Data analysis

After removing low quality reads, all clean data were used to perform K-mer analysis. Based on the results of the K-mer analysis, information on peak depth and the number of predicted best K-mer were obtained and used to estimate the size of the genome. Its relationship was expressed by using the following algorithm: Genome size =  $K\text{-mer\_num}/\text{peak\_depth}$ , where *K-mer\_num* is the total number of predicted best K-mer, and *peak\_depth* is the expected value of the K-mer depth. Also, the heterozygosity ratio and repeat sequence ratio were estimated following the description in [23], based on the K-mer analysis. K-mer analyses were performed using software GCE v1.0.0 [24] and KmerGenie v1.7039 [25], respectively. The clean reads were assembled into contigs in software SOAPdenovo v2.01 [26] with a K-mer of 21 by applying the *de Bruijn* graph structure. The paired-end information was then used to join the unique contigs into scaffolds.

The Perl script M*icro*S*atellite* (MISA, <http://pgrc.ipk-gatersleben.de/misa/>) was used to identify microsatellite motifs in the *de novo* draft genome. The search parameters were set for the detection of di-, tri-, tetra-, penta-, and hexanucleotide microsatellite motifs with a minimum of 6, 5, 5, 5, and 5 repeats, respectively. The microsatellite loci were subjected to primer design using Primer3 v2.3.7 software [27,28] with the standard parameters.

## Results and discussion

### Genome size prediction and sequence assembly

The experimental design, sequencing and analysis pipeline is shown in Figure 1. A total of 35.1 Gb raw data were generated by sequencing genome survey library with 350 bp inserts. The effective rate, error rate, Q20, Q30, and GC content of raw data was shown in Table 1 and Figure 2. A total of 34.8 Gb clean data were obtained after filtering and used for K-mer analysis. When employing KmerGenie, the predicted best K for K-mer analysis was 107 and the predicted genome size was about 812.86 Mb. Comparatively, when using GCE, the 21-mer frequency distribution derived from the sequencing reads is plotting in Figure 3; the peak of the 21-mer distribution was 38, and the total K-mer count was 29,998,886,801. As a result, the genome size was estimated as 796.25 Mb and the heterozygosity ratio and repeat sequence ratio were 0.17% and 39.65%, respectively. The development of NGS technology has provided researchers with an affordable way of addressing a wide range of questions, especially in non-model species such as *S. marmoratus* [11]. In addition, the K-mer method has been successfully applied for the estimation of genome size




Figure 1. Overview of the experimental design and analysis pipeline





Figure 2. Distribution figure of error rate, sequencing quality and GC content of raw data

using NGS reads without prior knowledge of the genome size [29]. Here, for the first time, we reported a genome survey of *S. marmoratus* using whole genome shotgun sequencing. The K-mer analyses suggested that the genome size is about 800 Mb, which is 87% of the size (920 Mb) previously estimated for *S. marmoratus* using flow cytometry [30].

Assembly was performed using 34.8-Gb Illumina PE clean reads. The length of contig N50 was 674 bp, and the Scaffold N50 was 4362 bp. The total length of scaffolds was 609.46 Mb. The GC content of scaffolds was 41.3%. The number of scaffolds >100 bp was 412,901 (98.99%) and >1 kb was 188,316 (45.15%) (Table 2). Information about the genome size of *S. marmoratus* from the present study may be useful for further genomic studies in this species.

## Identification and characteristic of microsatellite motifs in genome survey

From the 609,456,819 bp genome survey sequence, a total of 191,592 microsatellite motifs were identified, which included 140,801 microsatellite-containing sequences. However, only 67,846 sequences contained more than one



**Figure 3. K-mer (21-mer) analysis for estimating the genome size of *S. marmoratus***

The X-axis is depth and the Y-axis is the proportion that represents the frequency at that depth. Data produced from 350 bp insert library. The peak K-mer frequency was 38.

**Table 2 The result of assembly in *S. marmoratus* using 34.8-Gb Illumina clean data**

	Contigs		Scaffolds	
	Size (bp)	Number	Size (bp)	Number
N90	145	995,699	1117	179,431
N80	241	680,067	1877	128,640
N70	373	485,655	2589	94,551
N60	518	353,199	3413	69,208
N50	674	254,586	4362	49,699
Total size	583,830,195	–	609,456,819	–
GC content	41.39%	–	41.30%	–
Total number (> 100 bp)	1,467,661		412,901	
Total number (> 1 kb)	127,823		188,316	

microsatellite motifs, and 16,325 microsatellites were present in compound formation. Therefore, the microsatellite distribution frequency in this genome was estimated to be about 314.6 microsatellite per Mb. The motif types of microsatellites included 76.10% dinucleotide, 19.63% trinucleotide, 3.91% tetranucleotide, 0.36% pentanucleotide, and few hexanucleotide repeats (Figure 4A, Supplementary Table S1). The number of dinucleotide repeats was the highest, which was similar to previous studies on the distributions and characteristics of the microsatellites in *S. marmoratus* [22]. The frequency of repeats in most eukaryotes decreases exponentially with repeat length because mutation rates are higher in longer repeats [31]. Chen et al. [32] also reported that the number of repeats is inversely correlated with repeat length, and our present results confirmed this pattern. The relative abundances of specific repeat motifs were highly variable among the repeats. The frequency distribution range of microsatellite repeats ranged from 6 to 11 repeats for dinucleotide, from 5 to 8 repeats for trinucleotide, from 5 to 6 repeats for tetranucleotide. Of the dinucleotide repeats, the AC, TG, CA, and GT repeats were the first four repeats in abundance, accounting for 19.8% (28,853), 18.4% (26,797), 16.5% (24,093), and 14.3% (20,896), respectively (Figure 4B). Of the trinucleotide repeats, the GAG repeat was the most abundant, accounting for 5.4% (2030), whereas the ACG repeat was the least, accounting for 0.07% (25). In terms of the frequency of repeats, the 5-fold repeat was the most frequent of all trinucleotide repeats (Figure 4C). Of the tetranucleotide repeats, the ATAG repeat was the most abundant, accounting for 3.2% (239) (Figure 4D). However, only 5- and 6-fold repeats were identified and the 5-fold repeat was predominant of all tetranucleotide repeats. Compared with the results of Song et al. [22], in which the distributions and characteristics of the microsatellites in *S. marmoratus* were analyzed on the basis of 454 FLX pyrosequencing technique, our results showed high-efficiency in microsatellite loci identification. The number of microsatellites in the present study, as well as the kinds of microsatellite motifs, was much higher than previous study using 454 FLX pyrosequencing technique [22]. This difference might be due to the higher throughput of Illumina sequencing than 454 pyrosequencing.



## Author Contribution

T.X.G. conceived the study. S.Y.X., N.S. and S.J.X. collected the samples and extracted the genomic DNA. S.Y.X. performed genome assembly and bioinformatics analyses. S.Y.X. and T.X.G. wrote the original draft manuscript and all authors reviewed the manuscript.

## Ethics Approval

Ethical approval was not required for this study because no endangered or alive animals were involved. The specimen used in this study was caught by hook fishing and was dead when collected. All handling of *Sebastiscus marmoratus* specimens was conducted in strict accordance with Animal Care Quality Assurance in China and Zhejiang Ocean University.

## Abbreviations

GSS, Genome survey sequencing; NGS, next-generation sequencing; SSR, simple sequence repeat.

## References

- Moritz, C. (1994) Defining 'evolutionarily significant units' for conservation. *Trends Ecol. Evol.* **9**, 373–375, [https://doi.org/10.1016/0169-5347\(94\)90057-4](https://doi.org/10.1016/0169-5347(94)90057-4)
- Zane, L., Bargelloni, L. and Patarnello, T. (2002) Strategies for microsatellite isolation: a review. *Mol. Ecol.* **11**, 1–16, <https://doi.org/10.1046/j.0962-1083.2001.01418.x>
- Zhang, D.X. and Hewitt, G.M. (2003) Nuclear DNA analyses in genetic studies of populations: practice, problems and prospects. *Mol. Ecol.* **12**, 563–584, <https://doi.org/10.1046/j.1365-294X.2003.01773.x>
- Cheng, L., Liao, X., Yu, X. and Tong, J. (2007) Development of EST-SSRs by an efficient FIASCO-based strategy: a case study in rare minnow (*Gobiocypris Rarus*). *Anim. Biotechnol.* **18**, 143–152, <https://doi.org/10.1080/10495390601054980>
- Miller, M.R., Dunham, J.P., Amores, A., Cresko, W.A. and Johnson, E.A. (2007) Rapid and cost-effective polymorphism identification and genotyping using restriction site associated DNA (RAD) markers. *Genome Res.* **17**, 240–248, <https://doi.org/10.1101/gr.5681207>
- Triwitayakorn, K., Chatkulkawin, P., Kanjanawattanawong, S., Sraphet, S., Yoocha, T., Sangsrakru, D. et al. (2011) Transcriptome sequencing of *Hevea brasiliensis* for development of microsatellite markers and construction of a genetic linkage map. *DNA Res.* **18**, 471–482, <https://doi.org/10.1093/dnares/dsr034>
- Castoe, T.A., Poole, A.W., de Koning, A.P.J., Jones, K.L., Tomback, D.F., Oyler-McCance, S.J. et al. (2012) Rapid microsatellite identification from Illumina paired-end genomic sequencing in two birds and a snake. *PLoS One* **7**, e30953, <https://doi.org/10.1371/journal.pone.0030953>
- Capobianchi, M.R., Giombini, E. and Rozera, G. (2013) Next-generation sequencing technology in clinical virology. *Clin. Microbiol. Infect.* **19**, 15–22, <https://doi.org/10.1111/1469-0691.12056>
- Zhou, W., Hu, Y.Y., Sui, Z.H., Fu, F., Wang, J.G., Chang, L.P. et al. (2013) Genome survey sequencing and genetic background characterization of *Gracilariopsis lemaneiformis* (Rhodophyta) based on next-generation sequencing. *PLoS One* **8**, e69909, <https://doi.org/10.1371/journal.pone.0069909>
- Adelyna, M.A.N., Jung, H., Chand, V., Mather, P.B. and Azizah, M.N.S. (2016) A genome survey sequence (GSS) analysis and microsatellite marker development for Indian mackerel, *Rastrelliger kanagurta*, using Ion Torrent technology. *Meta Gene* **10**, 67–72, <https://doi.org/10.1016/j.mgene.2016.10.005>
- Motalebipour, E.Z., Kafkas, S., Khodaeiaminjan, M., Coban, N. and Gözel, H. (2016) Genome survey of pistachio (*Pistacia vera* L.) by next generation sequencing: development of novel SSR markers and genetic diversity in *Pistacia* species. *BMC Genomics* **17**, 998, <https://doi.org/10.1186/s12864-016-3359-x>
- Portis, E., Portis, F., Valente, L., Moglia, A., Barchi, L., Lanteri, S. et al. (2016) A genome-wide survey of the microsatellite content of the globe artichoke genome and the development of a web-based database. *PLoS One* **11**, e0162841, <https://doi.org/10.1371/journal.pone.0162841>
- Lu, X., Luan, S., Kong, J., Hu, L.Y., Mao, Y. and Zhong, S.P. (2017) Genome-wide mining, characterization, and development of microsatellite markers in *Marsipenaes japonicus* by genome survey sequencing. *Chin. J. Oceanol. Limnol.* **35**, 203–214, <https://doi.org/10.1007/s00343-016-5250-7>
- Fujita, H. and Kohda, M. (1998) Timing and sites of parturition of the viviparous scorpionfish, *Sebastiscus marmoratus*. *Environ. Biol. Fishes* **52**, 225–229, <https://doi.org/10.1023/A:1007471919373>
- Sun, D.Q., Shi, G., Liu, X.Z., Wang, R.X. and Xu, T.J. (2011) Genetic diversity and population structure of the marbled rockfish, *Sebastiscus marmoratus*, revealed by SSR markers. *J. Genet.* **90**, e21–e24
- Zhang, H. (2013) Molecular Phylogeography of Two Marine Ovoviviparous Fishes in Northwestern Pacific. Ph.D. Thesis, Ocean University of China, Qingdao, China (in Chinese)
- Xu, S.Y., Sun, D.R., Song, N., Gao, T.X., Han, Z.Q. and Shui, B.N. (2017) Local adaptation shapes pattern of mitochondrial population structure in *Sebastiscus marmoratus*. *Environ. Biol. Fishes* **100**, 763–774, <https://doi.org/10.1007/s10641-017-0602-5>
- Xu, T.J., Quan, X.Q., Sun, Y.N., Zhao, K.C. and Wang, R.X. (2010) A first set of polymorphic microsatellite loci from the marbled rockfish, *Sebastiscus marmoratus*. *Biochem. Genet.* **48**, 680–683, <https://doi.org/10.1007/s10528-010-9349-9>
- Yin, L.N., Zhang, H., Yanagimoto, T. and Gao, T.X. (2012) Isolation and characterization of nine polymorphic microsatellite markers of the marbled rockfish *Sebastiscus marmoratus* (Scorpaeniformes, Scorpaenidae). *Russian J. Genet.* **48**, 1264–1266, <https://doi.org/10.1134/S1022795412120174>
- Liu, H.B., Liu, S.F., Ye, J.B., Yuan, Y.J., Ding, S.X. and Zhuang, Z.M. (2014) Polymorphic microsatellite markers in the false kelpfish *Sebastiscus marmoratus*: isolation, characterization, and cross-species amplification. *Genet. Mol. Res.* **13**, 134–138, <https://doi.org/10.4238/2014.January.10.4>

- 21 Deng, H.W., Li, Z.B., Dai, G., Yuan, Y., Ning, Y.F., Shangguan, J.B. et al. (2015) Isolation of new polymorphic microsatellite markers from the marbled rockfish *Sebastes marmoratus*. *Genet. Mol. Res.* **14**, 758–762, <https://doi.org/10.4238/2015.January.30.19>
- 22 Song, N., Chen, M.Y., Gao, T.X. and Yanagimoto, T. (2017) Profile of candidate microsatellite markers in *Sebastes marmoratus* using 454 pyrosequencing. *Chin. J. Oceanol. Limnol.* **35**, 198–202, <https://doi.org/10.1007/s00343-016-5103-4>
- 23 Li, G., Song, L., Jin, C., Li, M., Gong, S.P. and Wang, Y.F. (2019) Genome survey and SSR analysis of *Apocynum venetum*. *Biosci. Rep.* **39**, BSR20190146, <https://doi.org/10.1042/BSR20190146>
- 24 Liu, B.H., Shi, Y.J., Yuan, J.Y., Hu, X.S., Zhang, H., Li, N. et al. (2013) Estimation of genomic characteristics by analyzing k-mer frequency in de novo genome projects. *arXiv*. **1308**, 2012, <https://arxiv.org/abs/1308.2012>
- 25 Chikhi, R. and Medvedev, P. (2014) Informed and automated k-mer size selection for genome assembly. *Bioinformatics* **30**, 31–37, <https://doi.org/10.1093/bioinformatics/btt310>
- 26 Luo, R.B., Liu, B.H., Xie, Y.L., Li, Z.Y., Huang, W.H., Yuan, J.Y. et al. (2012) SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. *GigaScience* **1**, 18, <https://doi.org/10.1186/2047-217X-1-18>
- 27 Koressaar, T. and Remm, M. (2007) Enhancements and modifications of primer design program Primer3. *Bioinformatics* **23**, 1289–1291, <https://doi.org/10.1093/bioinformatics/btm091>
- 28 Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B.C., Remm, M. et al. (2012) Primer3 - new capabilities and interfaces. *Nucleic Acids Res.* **40**, e115, <https://doi.org/10.1093/nar/gks596>
- 29 Lu, M., An, H. and Li, L. (2016) Genome survey sequencing for the characterization of the genetic background of *Rosa roxburghii* Tratt and leaf ascorbate metabolism genes. *PLoS One* **11**, e0147530, <https://doi.org/10.1371/journal.pone.0147530>
- 30 Ojima, Y. and Yamamoto, K. (1990) Cellular DNA contents of fishes determined by flow cytometry. *La Kromosomo II* **57**, 1871–1888
- 31 Katti, M.V., Ranjekar, P.K. and Gupta, V.S. (2001) Differential distribution of simple sequence repeats in eukaryotic genome sequences. *Mol. Biol. Evol.* **18**, 1161–1167, <https://doi.org/10.1093/oxfordjournals.molbev.a003903>
- 32 Chen, M., Tan, Z.Y., Zeng, G.M. and Peng, J. (2010) Comprehensive analysis of simple sequence repeats in pre-miRNAs. *Mol. Biol. Evol.* **27**, 2227–2232, <https://doi.org/10.1093/molbev/msq100>
- 33 Varshney, R.K., Graner, A. and Sorrells, M.E. (2005) Genic microsatellites markers in plants: features and application. *Trends Biotechnol.* **23**, 48–55, <https://doi.org/10.1016/j.tibtech.2004.11.005>
- 34 Queirós, J., Godinho, R., Lopes, S., Gortazar, C., de la Fuente, J. and Alves, P.C. (2015) Effect of microsatellite selection on individual and population genetic inferences: an empirical study using cross-specific and species-specific amplifications. *Mol. Ecol. Resour.* **15**, 747–760, <https://doi.org/10.1111/1755-0998.12349>

Table S1 Frequency of identified SSR motifs

Repeats	5	6	7	8	9	10	11	12	Total
AC	-	7749	5192	4967	7437	3450	58		28853
AG	-	3691	2063	1473	1287	1100	193		9807
AT	-	2457	1252	627	374	222	134	9	5075
CA	-	7026	4355	4029	5456	3103	70		24039
CG	-	37	14	1					52
CT	-	3453	2100	1354	1242	1128	165		9442
GA	-	2646	1585	1086	961	937	166		7381
GC	-	32	7	3	1				43
GT	-	5697	3700	3486	5172	2798	43		20896
TA	-	2249	1256	601	328	200	127	8	4769
TC	-	3111	1958	1208	1146	1053	172		8648
TG	-	8000	5144	4562	6175	2885	31		26797
AAC	327	135	124	13					599
AAG	355	169	137	11					672
AAT	814	512	384	6					1716
ACA	255	102	115	11					483
ACC	87	53	27	2					169
ACG	17	4	3	1					25
ACT	111	63	56	3					233
AGA	533	245	188	11					977
AGC	421	168	104	15					708
AGG	441	283	185	12					921
AGT	145	72	50	10					277
ATA	817	565	442	10					1834
ATC	171	95	118	15					399
ATG	177	104	133	11					425
ATT	788	454	357	9					1608
CAA	193	96	61	13					363
CAC	192	86	44	8					330
CAG	614	286	163	8					1071
CAT	274	160	180	10					624
CCA	259	142	64	6					471
CCG	85	25	15	4					129
CCT	780	554	289	4					1627
CGA	24	3	2						29
CGC	109	49	18	1					177
CGG	84	39	10	2					135
CGT	21	4	3						28
CTA	96	52	41	8					197
CTC	752	473	221	10					1456
CTG	477	248	151	13					889
CTT	340	197	197	9					743
GAA	304	139	170	16					629



GAC	38	24	4	3	69
GAG	1076	672	275	7	2030
GAT	185	94	118	16	413
GCA	396	173	113	10	692
GCC	99	42	11	6	158
GCG	91	29	14	4	138
GCT	508	215	166	10	899
GGA	613	416	223	4	1256
GGC	96	33	13	5	147
GGT	181	117	67	9	374
GTA	90	57	27	11	185
GTC	28	13	3	3	47
GTG	136	70	40	11	257
GTT	167	77	56	5	305
TAA	598	472	350	8	1428
TAC	183	95	65	7	350
TAG	137	67	42	7	253
TAT	625	397	345	5	1372
TCA	233	127	128	16	504
TCC	678	443	222	10	1353
TCG	23	6	1	1	31
TCT	399	226	153	11	789
TGA	212	120	149	15	496
TGC	362	168	91	3	624
TGG	160	87	33	12	292
TGT	291	132	117	7	547
TTA	730	458	377	12	1577
TTC	356	174	142	7	679
TTG	232	92	53	14	391
AAAC	49	19			68
AAAG	86	10			96
AAAT	183	6			189
AACA	26	5			31
AACC	8	2			10
AACG	1	2			3
AACT	7	9			16
AAGA	51	15			66
AAGC	3				3
AAGG	18	7			25
AAGT	4				4
AATA	133	7			140
AATC	41	10			51
AATG	44	15			59
AATT	29	5			34
ACAA	27	12			39
ACAG	90	10			100

ACAT	31	6	37
ACCA	9	2	11
ACCC	1		1
ACCG	3	2	5
ACCT	3		3
ACGC	6	10	16
ACGG		1	1
ACGT	1		1
ACTA	7	6	13
ACTC	3	5	8
ACTG	20	5	25
ACTT	2		2
AGAA	58	14	72
AGAC	92	10	102
AGAT	152	10	162
AGCC	6	4	10
AGCG	1		1
AGCT	7		7
AGGA	26	14	40
AGGC	5	4	9
AGGG	14	3	17
AGGT	4	1	5
AGTA	4	2	6
AGTC	10	7	17
AGTG	11	5	16
AGTT	17	6	23
ATAA	104	10	114
ATAC	35	8	43
ATAG	230	9	239
ATCA	39	16	55
ATCC	69	13	82
ATCG	1		1
ATCT	225	9	234
ATGA	49	13	62
ATGC	3	1	4
ATGG	58	12	70
ATGT	26	4	30
ATTA	34	8	42
ATTC	29	9	38
ATTG	39	7	46
ATTT	111	9	120
CAAA	17	10	27
CAAC	8	2	10
CAAG	1		1
CAAT	19	11	30
CACG	50	4	54

CACT	20	5	25
CAGA	103	10	113
CAGC	9	5	14
CAGG	10	6	16
CAGT	35	4	39
CATA	13	3	16
CATC	90	15	105
CATG	2	1	3
CATT	28	13	41
CCAA	7	6	13
CCAG	7	4	11
CCAT	59	13	72
CCCA	1		1
CCCT	6		6
CCGA	2	1	3
CCGT	3	2	5
CCTA	2	4	6
CCTC	18	1	19
CCTG	8	8	16
CCTT	15	3	18
CGAT	3		3
CGCA	9	7	16
CGGA	3	2	5
CGGT	1	2	3
CGTC	1		1
CGTG	6	9	15
CGTT	2		2
CTAA	12	1	13
CTAC	7	1	8
CTAT	124	12	136
CTCA	16	7	23
CTCC	16	1	17
CTGA	19	10	29
CTGC	12	2	14
CTGG	5	6	11
CTGT	122	9	131
CTTA	2	3	5
CTTC	25	8	33
CTTG	1	1	2
CTTT	42	13	55
GAAA	35	16	51
GAAG	26	5	31
GAAT	23	9	32
GACA	83	13	96
GACC	1	1	2
GACG	4	2	6

GACT	19	6	25
GAGC	2		2
GAGG	25	3	28
GAGT	8	6	14
GATA	130	11	141
GATC	1		1
GATG	72	13	85
GATT	13	11	24
GCAC	24	10	34
GCAG	12	7	19
GCAT	2		2
GCCA	6	1	7
GCCG	3		3
GCCT	9	4	13
GCGA	1		1
GCGG	1		1
GCGT	7	1	8
GCTA	6	2	8
GCTC	2		2
GCTG	8	5	13
GGAA	16	9	25
GGAC	8	1	9
GGAG	42	4	46
GGAT	65	11	76
GGCA	6	3	9
GGCT	9	5	14
GGGA	17	5	22
GGGT	1		1
GGTA	3	1	4
GGTC	1	1	2
GGTG	1	1	2
GGTT	4	1	5
GTAA	6	1	7
GTAG	8	2	10
GTAT	16	2	18
GTCA	16	4	20
GTCC	4	1	5
GTCG	4	1	5
GTCT	60	8	68
GTGA	14	4	18
GTGC	31	8	39
GTGG	1		1
GTTA	8	4	12
GTTC	1		1
GTTG	7		7
GTTT	21	7	28

TAAA	102	16	118
TAAC	5	4	9
TAAG	5	1	6
TAAT	12	4	16
TACA	9	6	15
TACC	4	1	5
TACT	4		4
TAGA	202	3	205
TAGC	11		11
TAGG	8	3	11
TAGT	12	7	19
TATC	177	12	189
TATG	27	8	35
TATT	125	8	133
TCAA	23	10	33
TCAC	11	9	20
TCAG	16	2	18
TCAT	33	10	43
TCCA	87	18	105
TCCC	11		11
TCCG	5	2	7
TCCT	36	7	43
TCGC	1		1
TCGG	2		2
TCTA	161	11	172
TCTG	121	5	126
TCTT	65	6	71
TGAA	47	14	61
TGAC	25	4	29
TGAG	14	4	18
TGAT	36	8	44
TGCA	3		3
TGCC	1		1
TGCG	8	8	16
TGCT	1		1
TGGA	61	13	74
TGGC	2	1	3
TGGG	3		3
TGGT	7	2	9
TGTA	30	9	39
TGTC	62	13	75
TGTT	52	11	63
TTAA	24	7	31
TTAC	8	4	12
TTAG	10	6	16
TTAT	100	11	111

TTCA	44	10	54
TTCC	20	11	31
TTCT	39	7	46
TTGA	20	8	28
TTGC	1		1
TTGG	3	1	4
TTGT	13	5	18
TTTA	141	12	153
TTTC	62	12	74
TTTG	31	3	34

---