

Research Article

ADSC-Exos containing MALAT1 promotes wound healing by targeting miR-124 through activating Wnt/ β -catenin pathway

 Lin He^{1,*}, Chan Zhu^{2,*}, Jing Jia¹, Xiao-Yan Hao¹, Xue-Yuan Yu¹, Xiang-Yu Liu¹ and Mao-Guo Shu¹

¹Department of Plastic, Aesthetic and Maxillofacial Surgery, The First Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710061, P.R. China; ²Department of Burns and Cutaneous Surgery, Xijing Hospital, The Fourth Military Medical University, Xi'an 710032, P.R. China

Correspondence: Lin He (helin637@163.com)



Cutaneous wound is a soft tissue injury that is difficult to heal during aging. It has been demonstrated that adipose-derived stem cells (ADSCs) and its secreted exosomes exert crucial functions in cutaneous wound healing. The present study aimed to elucidate the mechanism of exosomes derived from ADSCs (ADSC-Exos) containing MALAT1 in wound healing. ADSCs were isolated from human normal subcutaneous adipose tissues and identified by flow cytometry analysis. Exosomes were extracted from ADSC supernatants and MALAT1 expression was determined using qRT-PCR analysis. HaCaT and HDF cells were exposed to hydrogen peroxide (H₂O₂) for simulating the skin lesion model. Subsequently, CCK-8, flow cytometry, wound healing and transwell assays were employed to validate the role of ADSC-Exos containing MALAT1 in the skin lesion model. Besides, cells were transfected with sh-MALAT1 to verify the protective role of MALAT1 in wound healing. The binding relationship between MALAT1 and miR-124 were measured by dual-luciferase reporter assay. ADSC-Exos promoted cell proliferation, migration, and inhibited cell apoptosis of HaCaT and HDF cells impaired by H₂O₂. However, the depletion of MALAT1 in ADSC-Exos lose these protective effects on HaCaT and HDF cells. Moreover, miR-124 was identified to be a target of MALAT1. Furthermore, ADSC-Exos containing MALAT1 could mediate H₂O₂-induced wound healing by targeting miR-124 and activating Wnt/ β -catenin pathway. ADSC-Exos containing MALAT1 play a positive role in cutaneous wound healing possibly via targeting miR-124 through activating the Wnt/ β -catenin pathway, which may provide novel insights into the therapeutic target for cutaneous wound healing.

Introduction

Cutaneous wound healing is a dynamic process and involves four precisely integrated and overlapping phases, including hemostasis, inflammation, proliferation and tissue remodeling [1,2]. The wound has a long healing cycle, the nonhealing cutaneous wounds without effective therapies can lead to severe clinical burden [3]. Ineffective skin wound healing is an important source of morbidity and mortality. Importantly, the risk of chronic skin wounds failing to heal increases as the age increases [4]. In the process of skin wound healing, the integration of skin cell differentiation, migration, proliferation and apoptosis plays a crucial role in skin tissue repair [5]. The wound after healing has the characteristic of regenerative epithelialization, which is related to two basic functions of keratinocytes, namely proliferation and migration [6]. Some growth factors, such as transforming growth factor β (TGF- β) and epidermal growth factor (EGF), can stimulate cell proliferation, differentiation and migration. Many signaling pathways play crucial role during cutaneous wound repair: the Wnt/ β -catenin, Notch, Hedgehog and various growth factor/cytokine pathways [7]. The functions of these growth factors and signaling pathways in clinical

*These are co-first authors.

Received: 10 October 2019

Revised: 10 April 2020

Accepted: 20 April 2020

Accepted Manuscript online:

28 April 2020

Version of Record published:

11 May 2020

research confirm their role in accelerating wound healing. Though certain progress is obtained in wound healing, the particular mechanism of cutaneous wound healing has not been fully clarified.

Mesenchymal stem cells (MSCs) are pluripotent stem cells with differentiation abilities, and have been widely used in the field of regenerative medicine because of their important role in enhancing the ability of various tissues to regenerate [8,9]. Numerous studies have reported the significant role of MSCs in neovascularization of ischemic tissue, including wound healing [10,11]. Exosomes are small multivesicular intraluminal vesicles ranging from 30 to 100 nanometers in diameter [12]. Exosomes are important paracrine mediators between MSCs and target cells, enabling cell-to-cell communication by delivering RNA and proteins to target cells [13,14]. Li et al. [15] reported that exosomes from adipose-derived stem cells (ADSCs) can potentially promote wound healing. Dalirfardouei et al. [16] suggested that the menstrual blood-derived MSCs-derived exosomes effectively ameliorated cutaneous nonhealing wounds. Ma et al.'s [17] findings also revealed that exosomes derived from ADSCs (ADSC-Exos) play a positive role in cutaneous wound healing possibly via Wnt/ β -catenin signaling. MALAT1 may act as a transcriptional regulator for numerous genes, including some genes involved in cancer metastasis and cell migration, and it is involved in cell cycle regulation [18]. CD63 mediates signal transduction events that play a role in the regulation of cell development, activation, growth and motility [19]. CD9 functions in many cellular processes including differentiation, adhesion and signal transduction and expression of this gene plays a critical role in the suppression of cancer cell motility and metastasis [20]. However, the molecular mechanism of ADSCs in wound healing has not been fully investigated.

In the present study, we isolated exosomes from ADSCs and established a skin lesion model via exposure of HaCaT and HDF cells to hydrogen peroxide (H_2O_2), and then investigated the wound healing properties of exosomes derived from ADSCs. We demonstrated that ADSCs-Exos containing MALAT1 significantly attenuated H_2O_2 -induced the suppression of cell proliferation, migration and the promotion of apoptosis via targeting miR-124 through Wnt/ β -catenin pathway. Collectively, our data implied that MALAT1 may play critical regulatory roles in promoting cutaneous wound healing.

Materials and methods

Isolation and culture of ADSCs

Human normal subcutaneous adipose tissues (obtained in healthy people undergoing selective liposuction or undergoing surgical plastic surgery) were acquired from the First Affiliated Hospital of Xi'an Jiaotong University. The Research Ethics Committee of the First Affiliated Hospital of Xi'an Jiaotong University approved the study protocol, and informed consent was obtained from all the donors of adipose tissue. Then the adipose tissues were digested by collagenase type 1 for 60 min and was terminated using DME/F12 complete media. The cell-debris pellet was obtained by centrifugation at 1000 rpm for 5 min. The cells were cultured in a 5% CO_2 humidified atmosphere at 37°C, and the medium was changed every 3 days. Upon reaching 80% confluence, the cells were defined as passage 1 and re-plated until the fourth passage for the following experiments.

Flow cytometry analysis

The fourth-passage human ADSCs were tested by flow cytometry analysis. Briefly, the adherent cells were harvested by trypsinization, centrifuged at 800 rpm for 6 min, washed with sterile phosphate-buffered saline and resuspended. Cells were stained with specific fluorescein isothiocyanate-conjugated (FITC) monoclonal antibodies against CD29, CD44, CD34 and CD31 for 45 min at 4°C. For the negative control, an irrelevant antibody of the same isotype was used. Finally, the cells were analyzed using an FACS Calibur cytometer [21].

Exosomes isolation and identification

Using exosome isolation reagent (RiboBio, Guangzhou, China), exosomes were extracted from ADSCs supernatants without cell-debris in accordance with the manufacturer's guidelines. Subsequently, observation of isolated exosomes was monitored by means of transmission electron microscopy (TEM; JEOL Ltd., JEM2010-HT; Tokyo, Japan). The ADSCs-Exos were characterized by nanoparticle tracking analysis (NTA) and Western blotting. In the NTA assay, the size, distribution and the number of particles in the ADSCs-Exo were evaluated using a nanoparticle tracking analyzer (v3.1, Malvern Instruments, Ltd., Worcestershire, U.K.) [22]. Total RNA and protein were extracted using TRIzol-LS (Invitrogen, Carlsbad, CA, U.S.A.) and Exosomal Protein Extraction kit (Invitrogen), according to the manufacturer's protocol, respectively. The ADSCs-Exo were analyzed by Western blotting using the following primary antibodies: anti-CD9 (1:1000, Abcam), anti-CD63 (1:1000, Abcam). Final exosomes were obtained and stored at $-80^\circ C$ for use for the following study.

Cell culture and cell transfection

HaCaT cells were purchased from the American Type Culture Collection (ATCC, Manassas, U.S.A.) and HDF cells were obtained from ScienCell. Cells were cultured in Dulbecco's modified Eagle's medium (DMEM; HyClone, Logan) and 10% fetal bovine serum without exosomes (Gibco, Carlsbad, CA) and maintained in a humidified incubator at 37°C with 5% CO₂.

For MALAT1 knockdown, MALAT1-targeting shRNAs (GenePharma, Shanghai, China) was synthesized to knockdown the expression of MALAT1 (sh-MALAT1). A negative control (GenePharma) was named as shNC. Anti-miR-124 was utilized to change the expression level of miR-124 in HaCaT and HDF cells. One control miRNAs (miR-NC) was used as negative controls for anti-miR-124. Cell transfection was then performed using Lipofectamine 2000 (Invitrogen) according to the manufacturer's instructions.

MTT assay

MTT assay was used to evaluate the survival of HaCaT and HDF cells with different H₂O₂ treatment (0, 100, 500, 1000, 1500 and 2000 μM). The details of MTT assay was performed as previously described [23].

RNA isolation and real-time PCR

Total RNA was extracted with TRIzol Reagent (Invitrogen), followed by cDNA synthesis using a TransScript All-in-One First-Strand cDNA Synthesis SuperMix (Transgen Biotech, Beijing, China), was performed as described previously [24]. PCR was performed using a Bio-Rad PCR instrument (Bio-Rad, Hercules, CA, U.S.A.) with 2× Taq PCR Master Mix (Solarbio, Beijing, China) following the manufacturer's instructions. The fold changes were calculated by means of relative quantification ($2^{-\Delta\Delta C_t}$ method). PCR primers are described as below:

MALAT1: forward 5'-GGGTGTTTACGTAGACCAGAACC-3' and reverse 5'-CTTCCAAAAGCCTTCTGCCTTAG-3'; miR-124: forward 5'-TCGTTAAGGCACGCGGTG-3' and reverse 5'-GTGCAGGGTCCGAGGT-3'; U6: forward 5'-CTCGCTTCGCAGCACACA-3' and reverse 5'-AACGCTTCACGAATTTGCGT-3'; GAPDH forward 5'-GCACCGTCAAGGCTGAGAAC-3' and reverse 5'-ATGGTGGTGAAGACGCCAGT-3'.

CCK-8 assay

The capacity of cell proliferation was assessed using CCK-8 (Beyotime Biotechnology). Cells were planted into 96-well plates (5000 cells/well) containing culture medium. Subsequently, cells were exposed to different concentrations of H₂O₂ at indicated times. CCK-8 assay was carried out by adding 10 μl of CCK-8 reagent into each well. After incubated in an incubator for another 2 h, cell proliferation was measured by testing the absorbance at 450 nm using a microplate reader (Bio-Rad Laboratories, Inc., Hercules, CA, U.S.A.).

Migration assay

The migratory properties of HaCaT and HDF cells were tested by two methods. (1) For the scratch wound healing assay, HaCaT and HDF cells were seeded into plastic six-well plates at the density of 5×10^5 cells per well and cultured for 12 h. Cells were treated with 500 μM of H₂O₂ for 4 h when they reached a confluence of 80%. After discarding the culture medium, uniform scratch wounds were scraped by a sterile pipette tip. Each well was washed with PBS, and then supplemented with basal DMEM containing H₂O₂, H₂O₂+ADSC-Exo, H₂O₂+ADSC-Exo-shNC, H₂O₂+ADSC-Exo-shMALAT1, H₂O₂+ADSC-Exo-shMALAT1+anti-miR-124, respectively. Images for each scratch were observed by microscope and captured at 0 and 24 h after scratching. (2) For the transwell assay, transwell chambers (8 μM, Corning Incorporated, Corning, NY) without Matrigel were applied following the manufacturer's protocol. Cells were treated with H₂O₂, H₂O₂+ADSC-Exo, H₂O₂+ADSC-Exo-shNC, H₂O₂+ADSC-Exo-shMALAT1, H₂O₂+ADSC-Exo-shMALAT1+anti-miR-124, respectively. Subsequently, the harvested cells were suspended in 200 μl of serum-free media and were added into the upper chamber, and the lower chamber was filled with normal growth media. After 24 h incubation, the migrated cells were fixed with 4% formaldehyde and stained with 0.1% Crystal Violet solution [17].

Dual-luciferase reporter assay

The 3'UTR of MALAT1 gene containing the predicated binding sites for miR-124 were amplified using PCR. The fragment was inserted into the multiple cloning sites in the pMIR-REPORT luciferase miRNA expression reporter vector (Ambion, Austin, U.S.A.). Then, HEK293T cells were co-transfected with 0.1 μg of luciferase reporter vectors comprising wild-type or mutant type of 3'UTR of MALAT1 and either miR-124 mimic or miR-control by Lipofectamine 2000 (Invitrogen, Carlsbad, U.S.A.). Relative luciferase activity was calculated by normalizing the firefly




Figure 1. Isolation and characterization of ADSCs and ADSC-Exos

(A) The expression of biomarkers (CD34, CD31, CD29 and CD44) were detected by FACS. (B) The exosomes' marker proteins (CD63, TSG101 and CD9) detected by Western blot. (C) The expression level of MALAT1 was measured using qRT-PCR assay. *** $P < 0.001$.

luminescence to the *Renilla* luminescence using the Dual-Luciferase Reporter Assay System (Promega, Madison, WI, U.S.A.) according to the manufacturer's instructions at 48 h post-transfection.

Statistical analysis

Experiments were independently performed at least three times. All data were presented as mean \pm standard deviation (SD) and the differences among multiple groups were analyzed by Student's *t* test or one-way analysis of variance (ANOVA). The differences were considered to be statistically significant as P -value < 0.05 .

Results

Isolation and characterization of ADSCs and ADSC-Exos

After primary isolation, ADSCs were cultured for 24 h. Subsequently, we observed that most of the adherent cells were in spindle-like shape during cell culture using a microscope ($\times 40$ magnification) (Supplementary Figure S1). Besides, flow cytometry analysis was performed to measure the relevant biomarkers. As shown in Figure 1A, CD29 and CD44 (MSC markers) were highly expressed in ADSCs at passage 4, while the expression level of CD34 and CD31 (hematopoietic cell markers) were not expressed in the ADSCs (Figure 1B). In addition, exosomes purified from ADSCs culture supernatants were characterized by TEM, and the results showed that exosomes were round membrane-bound vesicles with 30–100 nm diameter (Supplementary Figure S2). Meanwhile, CD9, TSG101 and CD63, which seemed to be the exosomes marker protein, were also detected in the exosomes as expected (Figure




Figure 2. Construction of the skin lesion model *in vitro*

(A) Cell viability assessed by MTT assay. (B) Apoptotic rate of HaCaT and HDF cells exposed with different concentrations of H_2O_2 examined by flow cytometry assay. (C) The percentage of apoptosis cell for HaCaT and HDF cells. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

1B). Furthermore, the gene level of MALAT1 was highly expressed in ADSC-exos compared with ADSCs (Figure 1C).

Construction of the skin lesion model

To simulate the skin lesion model *in vitro*, HaCaT and HDF cells were treated with H_2O_2 at different concentrations of 0, 100, 500, 1000, 1500 and 2000 μM . As shown in Figure 2A, cell viability of HaCaT and HDF cells was significantly inhibited by the application of H_2O_2 in a dose-dependent manner, indicating that H_2O_2 could induce skin lesions. Consistently, the results obtained from flow cytometry assay (Figure 2B,C) showed that H_2O_2 also promoted the apoptosis of HaCaT and HDF cells, compared with that of control. Besides, as shown in Western blot analysis (Supplementary Figure S3), the results revealed that the increased expression of Caspase-3/Bax and the decreased expression of Bcl-2 were detected, with the enhanced concentrations of H_2O_2 .

ADSC-Exos attenuate the suppression of cell proliferation, migration and the promotion of cell apoptosis in HaCaT and HDF cells impaired by H_2O_2

To explore the biological functions of ADSC-Exos in the cutaneous wound healing, the following experiments were employed. As shown in Supplementary Figure S4, cell proliferation was obviously repressed by H_2O_2 , while the suppressive effects were reversed after the addition of ADSC-Exos. Meanwhile, flow cytometry also presented that ADSC-Exos significantly reduced the apoptotic rate of HaCaT and HDF cells induced by H_2O_2 (Figure 3A,B). Moreover, the capacity of cell migration was also determined using the scratch wound healing (Supplementary Figure S5) and transwell assays (Figure 3C,D). As expected, the migratory ability of HaCaT and HDF cells were impaired by H_2O_2 , but the migratory ability was increased after adding the ADSC-Exos. These findings implied the protective role of ADSC-Exos in the cutaneous wound healing.




Figure 3. Effects of ADSC-Exos on cell proliferation, apoptosis and migration of HaCaT cells impaired by H₂O₂

(A) Apoptosis of HaCaT and HDF cells was evaluated by flow cytometry assay. (B) The percentage of apoptosis cell for HaCaT and HDF cells. (C,D) The capacity of cell migration in HaCaT and HDF cells was analyzed by transwell assay. **P*<0.05, ***P*<0.01, ****P*<0.001.




Figure 4. ADSC-Exos knocking down MALAT1 lose protective effects on HaCaT and HDF cells

(A) The expression level of MALAT1 was measured by qRT-PCR assay. (B) Proliferation of HaCaT and HDF cells treated with H₂O₂, H₂O₂+ADSC-Exo, H₂O₂+ADSC-Exo-shNC or H₂O₂+ADSC-Exo-shMALAT1 were evaluated by CCK-8 assay. (C, D) Migration of HaCaT and HDF cells treated with H₂O₂, H₂O₂+ADSC-Exo, H₂O₂+ADSC-Exo-shNC or H₂O₂+ADSC-Exo-shMALAT1 were analyzed by Transwell assay. (E) The expression level of miR-124 was performed using qRT-PCR analysis. **P*<0.05, ***P*<0.01, ****P*<0.001.

ADSC-Exos knocking down MALAT1 lose the protective effects on HaCaT and HDF cells impaired by H₂O₂

To further validate whether ADSC-Exo containing MALAT1 play a role in mediating skin lesion model, we performed a pretreatment of ADSCs with MALAT1 knockdown, followed by collection of conditioned supernatants for exosomal extraction, namely ADSC-Exo-shMALAT1. As shown in Figure 4A, qRT-PCR assay confirmed that the expression level of MALAT1 was down-regulated after MALAT1 knockdown (for HaCaT approximately 40%, for HDF approximately 30%). As shown in Figure 4B, ADSC-Exos dramatically attenuated the impairment of H₂O₂ treatment, on cell proliferation in HaCaT and HDF cells. However, the positive effect of ADSC-Exos was significantly inhibited after knocking down MALAT1 (Figure 4B). Similarly, flow cytometry assay also showed dramatically decreased apoptosis of HaCaT and HDF cells after adding ADSC-Exos and obvious increased apoptosis after MALAT1 knockdown (Supplementary Figure S6). Additionally, the results of scratch wound healing assay also implied that the migratory ability of HaCaT and HDF cells were decreased by H₂O₂, but were increased after adding the ADSC-Exos. However, the promotion of cell migration induced by ADSC-Exos was reversed within MALAT1 knockdown (Supplementary Figure S7). Similar results were obtained from transwell assay (Figure 4C,D). Furthermore, the expression level of




Figure 5. MALAT1 acts as a sponge of miR-124

(A) Analysis of binding sites of MALAT1 and miR-124 using bioinformatics software. (B) The wild-type and mutant-type of the complementary binding sequences between MALAT1 and miR-124. (C) The binding relationship between MALAT1 and miR-124 was employed using dual-luciferase reporter assay. (D) qRT-PCR was used to detect the expression of miR-124 after transfection with sh-MALAT1. ** $P < 0.01$.

miR-124 detected by qRT-PCR was decreased after adding ADSC-Exos, but was increased after MALAT1 knock-down (Figure 4E). Taken together, ADSCs-Exo containing MALAT1 exert a protective role in the cutaneous wound healing.

MALAT1 acts as a sponge of miR-124

We used bioinformatics software to predict the binding sites of MALAT1 and miR-124, which was shown in Figure 5A. The complementary nucleotides between miR-124 and the seed region in MALAT1 reporter plasmids (WT) of 'GUGCCUUG' were replaced by 'CACGGAU' to construct MALAT1 mutant (Figure 5B). To identify whether MALAT1 directly targets miR-124, luciferase activity was assessed after co-transfection. As shown in Figure 5C, miR-124 mimic markedly decreased the luciferase activity of HEK-293T transfected with plasmid containing MALAT1-WT but not MALAT1-Mut. We also utilized qRT-PCR to study the expression level of miR-124 in shNC and shMALAT1 groups, which was shown in Figure 5D. There was higher expression level of miR-124 in the shMALAT1 group compared with shNC group.

ADSC-Exos containing MALAT1 mediates H₂O₂-induced wound healing by targeting miR-124 through activating the Wnt/ β -catenin pathway

It has been demonstrated that Wnt/ β -catenin signaling is related with wound healing [25], following that, whether Wnt/ β -catenin signaling is also involved in ADSC-Exo containing MALAT1 in wound healing was further explored. As shown in Figure 6A, the expression level of miR-124 was markedly decreased in anti-miR-124 group compared with that of control. Subsequently, the results of CCK-8 assay presented that ADSC-Exos relieved the inhibition of H₂O₂ on proliferation ability of HaCaT and HDF cells, while MALAT1 knockdown recovered the impairment of H₂O₂ on cell viability. However, the capacity of cell proliferation was recovered after silencing of miR-124 (Figure 6B). Similarly, ADSC-Exos relieved the apoptosis induced by H₂O₂ in HaCaT and HDF cells, but the effects were




Figure 6. ADSC-Exos containing MALAT1 mediates H₂O₂ induced wound healing by targeting miR-124 through activating Wnt/ β -catenin pathway

(A) The expression level of miR-124 was detected by qRT-PCR after transfected with anti-miR-124. (B) CCK-8 assay was performed to assess the cell proliferation of HaCaT and HDF cells treated with H₂O₂, H₂O₂+ADSC-Exo, H₂O₂+ADSC-Exo-shNC, H₂O₂+ADSC-Exo-shMALAT1, H₂O₂+ADSC-Exo-shMALAT1+anti-miR-124. (C) Flow cytometry was subjected to evaluate cell apoptosis of HaCaT and HDF cells treated with H₂O₂, H₂O₂+ADSC-Exo, H₂O₂+ADSC-Exo-shNC, H₂O₂+ADSC-Exo-shMALAT1, H₂O₂+ADSC-Exo-shMALAT1+anti-miR-124. (D) Migration of HaCaT and HDF cells treated with H₂O₂, H₂O₂+ADSC-Exo, H₂O₂+ADSC-Exo-shNC, H₂O₂+ADSC-Exo-shMALAT1, H₂O₂+ADSC-Exo-shMALAT1+anti-miR-124 were analyzed by the scratch wound healing assay. (E) The expression of Wnt/ β -catenin signals were determined by western blot. (F) Cell proliferation of HaCaT and HDF cells treated with FH535 were evaluated by CCK-8 assay. (G) Wnt/ β -catenin signal pathway of HaCaT and HDF cells treated with H₂O₂, H₂O₂+ADSC-Exo or H₂O₂+ADSC-Exo+FH535 were monitored by western blot. **P*<0.05, ***P*<0.01, ****P*<0.001.

blocked by MALAT1 knockdown. Moreover, the apoptosis was decreased after silencing the expression of miR-124 (Figure 6C). Besides, the inhibitory effect of H₂O₂ on migration ability of HaCaT and HDF cells was restrained after adding ADSC-Exos, while down-regulation of MALAT1 impaired the protection of ADSC-Exos on the migration ability. However, the effects were recovered by silencing of miR-124 (Figure 6D). In addition, Wnt/ β -catenin signaling was detected using Western blot and the results showed that Wnt/ β -catenin signaling was activated after adding ADSC-Exos, and MALAT1 knockdown further caused the inactivation of Wnt/ β -catenin signaling pathway. However, silencing of miR-124 regained the promotion of Wnt/ β -catenin signaling (Figure 6E). Furthermore, FH535, an inhibitor of Wnt/ β -catenin signaling, notably impaired the protection of ADSC-Exos on cell proliferation and migration (Figure 6F and Supplementary Figure S8). As expected, compared with control group, FH535 significantly blocked the activation of Wnt/ β -catenin signaling pathway (Figure 6G).

Discussion

Although medical technology has made great progress in promoting wound healing over the past decades, scar formation during healing remains a challenging clinical problem [26]. Cell therapy has shown potential to enhance skin wound healing and decrease fibrosis after wounding [27]. The potential use of MSCs has been a hot topic in tissue repair studies [28]. ADSCs have the characteristics of pluripotent differentiation and extended proliferation ability, and can be harvested in large quantities [29]. Previous studies have indicated that ADSCs secrete several trophic factors and have a degree of immune privilege because of their ability to suppress T-cell-mediated responses that cause tissue rejection [30]. ADSCs have been successfully used in the treatment of soft tissue defects, scars and burns by accelerating and improving the quality of the wound healing process. ADSCs promote angiogenesis, epithelial cell migration, growth factor secretion and differentiation into multiple lineages [31], thus enhancing wound healing and reducing scar formation. However, the exact mechanism still remains elusive. In the present study, we exposed HaCaT and HDF cells to H₂O₂ to simulate a skin lesion model. The results of increased apoptosis, increased Bax expression and decreased Bcl-2 expression confirmed the validity of the model.

Exosomes are released from cells and participate in many biological and pathological processes. Since exosomes can carry messenger RNA, microRNAs and proteins, which can be transferred to target cells, the novel role of exosomes as cell communication bodies has been verified in many researches. The promotive role of exosomes in the proliferation, migration and angiogenesis process has been demonstrated. There is increasing evidence showing that the ability of ADSCs to stimulate skin fibroblast proliferation during wound healing might be due to the presence of growth factors secreted by ADSCs, especially exosomes. Therefore, we aimed to investigate whether ADSC-Exos exert protective functions in wound healing. In the present study, the existence of exosomes derived from ADSC was evidenced by CD63, TSG101 and CD9 expression. Interestingly, we observed that exosomes derived from ADSCs significantly promoted proliferation, migration and inhibited apoptosis of HaCaT and HDF cells induced by H₂O₂, these findings are consistent with data reported by other researchers, who have indicated that human ADSCs accelerate the repair process in various adult tissues and improve the quality of repair via exosomes [32–34]. Further investigation revealed that the positive effects of ADSC-Exos were blocked by MALAT1 knockdown. To explore the molecular mechanism of ADSC-Exos in wound healing, the following researches were conducted.

MALAT1 was one of the first identified lncRNAs associated with human diseases, which was originally described to be associated with metastasis of lung cancer [35]. It has been reported that MALAT1 drives tumorigenesis through the promotion of tumor cell proliferation. Moreover, overexpression of MALAT1 results in increased cell migration *in vitro* [36]. Depletion of MALAT1, on the other hand, inhibits cell motility *in vitro* and significantly limits metastasis formation in mouse cancer models [37]. Besides, it has also been reported that MALAT1 is involved in

diabetes-induced microvascular dysfunction and regulates retinal endothelial cell proliferation, migration and tube formation [38]. Previous study reported that MALAT1 increased cell migration via modulating miR-140 expression in a uveal melanoma cell line [39]. MALAT1 can also up-regulate the expression of miR-22-3p target genes CXCR2 and Akt [40]. However, the exact role of MALAT1 on wound healing has not been clearly identified despite its extensive investigation in the cancer. Herein, we investigated the potential role of lncRNA MALAT1 in wound healing via MALAT1 knockdown in ADSC-Exos. The expression level of miR-124 was significantly increased after MALAT1 knockdown. Dual-luciferase reporter assay also verified the direct binding between MALAT1 and miR-124.

As previously described, Zhang et al. [41] found that miR-124 inhibited keratinocyte proliferation, collagen biosynthesis and activation of Wnt/ β -catenin by targeting SERP1. Yang et al. [42] reported that miR-124 inhibited proliferation, migration and invasion of malignant melanoma (A375) cells. The function of Wnt/ β -catenin signaling pathway has been widely studied; it is involved not only in cell proliferation and cell cycle, but also in wound healing [43,44]. In addition, the Wnt/ β -catenin pathway has been shown to regulate keratinocyte proliferation and apoptosis in psoriasis lesions. Therefore, we investigated the involvement of Wnt/ β -catenin pathway in wound healing. In our results, MALAT1 knockdown caused the inactivation of Wnt/ β -catenin signaling pathway. However, silencing of miR-124 recovered the Wnt/ β -catenin signaling pathway. ADSC-Exos induced cell viability was inhibited after adding FH535, an inhibitor of Wnt/ β -catenin signaling pathway. Similarly, the migratory ability of HaCaT and HDF cells was also repressed by FH535 treatment. These results suggested that Wnt/ β -catenin pathway was involved in the wound healing, which was activated by MALAT1.

Finally, we have shown that ADSC-Exos containing MALAT1 significantly promote cell proliferation, migration and inhibit cell apoptosis in the skin lesion models. miR-124 is a target gene of MALAT1 and its downexpression could attenuate the protective effects of ADSC-Exos on cutaneous cells. Moreover, MALAT1 significantly activates the Wnt/ β -catenin pathway. These findings implied that ADSC-Exos containing MALAT1 mediates H₂O₂-induced wound healing by targeting miR-124 and activating the Wnt/ β -catenin pathway. Nevertheless, the effect and underlying mechanism of MALAT1 and miR-124 on the development of wound healing are still in their infancy. Further research is needed to better understand the mechanism whereby MALAT1 regulates the progression of wound healing before MALAT1-based therapeutics can be taken into clinical practice.

As a pilot study, we are currently screening the mechanism of ADSC-Exos containing MALAT1 in wound healing by using ADSCs and its secreted exosomes only at the cellular level. Our future *in vivo* molecular intervention experiments will extend our current findings.

Competing Interests

The authors declare that there are no competing interests associated with the manuscript.

Funding

This work was supported by the Shannxi Key R&D Plan — Social Development Area [grant number 2019SF-106].

Author Contribution

Guarantor of integrity of the entire study: Lin He. **Study concepts:** Lin He. **Study design:** Lin He. **Definition of intellectual content:** Lin He. **Literature research:** Jing Jia. **Clinical studies:** Lin He, Xue-Yuan Yu and Xiang-Yu Liu. **Experimental studies:** Lin He, Chan Zhu and Jing Jia. **Data acquisition:** Chan Zhu; **Data analysis:** Xiao-Yan Hao. **Statistical analysis:** Chan Zhu. **Manuscript preparation:** Lin He. **Manuscript editing:** Lin He. **Manuscript review:** Mao-Guo Shu.

Ethics Approval

The Research Ethics Committee of the First Affiliated Hospital of Xi'an Jiaotong University approved the study protocol, and informed consent was obtained from all the donors of adipose tissue. The research has been carried out in accordance with the World Medical Association Declaration of Helsinki (2013 version), and that all subjects provided written informed consent.


Abbreviations

ADSC, adipose-derived stem cell; ADSC-Exos, exosomes derived from ADSC; CCK-8, cell counting kit-8; DMEM, Dulbecco's modified Eagle's medium; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; H₂O₂, hydrogen peroxide; MALAT1, metastasis associated lung adenocarcinoma transcript 1; MSC, mesenchymal stem cell; NTA, nanoparticle tracking analysis; qRT-PCR, quantitative real-time polymerase chain reaction; SERP1, stress-associated endoplasmic reticulum protein 1.

References


- 1 Rodriguez-Menocal, L., Salgado, M., Ford, D. and Van Badiavas, E. (2012) Stimulation of skin and wound fibroblast migration by mesenchymal stem cells derived from normal donors and chronic wound patients. *Stem Cells Transl. Med.* **1**, 221–229, <https://doi.org/10.5966/sctm.2011-0029>
- 2 Guo, S. and Di Pietro, L.A. (2010) Factors affecting wound healing. *J. Dent. Res.* **89**, 219–229, <https://doi.org/10.1177/0022034509359125>
- 3 Zhao, P., Sui, B.D., Liu, N., Lv, Y.J., Zheng, C.X., Lu, Y.B. et al. (2017) Anti-aging pharmacology in cutaneous wound healing: effects of metformin, resveratrol, and rapamycin by local application. *Aging Cell* **16**, 1083–1093, <https://doi.org/10.1111/ace1.12635>
- 4 Sgonc, R. and Gruber, J. (2013) Age-related aspects of cutaneous wound healing: a mini-review. *Gerontology* **59**, 159–164, <https://doi.org/10.1159/000342344>
- 5 Lewis, C.J., Mardaryev, A.N., Poterlowicz, K., Sharova, T.Y., Aziz, A., Sharpe, D.T. et al. (2014) Bone morphogenetic protein signaling suppresses wound-induced skin repair by inhibiting keratinocyte proliferation and migration. *J. Invest. Dermatol.* **134**, 827–837, <https://doi.org/10.1038/jid.2013.419>
- 6 Yang, L., Zheng, Z., Zhou, Q., Bai, X., Fan, L., Yang, C. et al. (2017) miR-155 promotes cutaneous wound healing through enhanced keratinocytes migration by MMP-2. *J. Mol. Histol.* **48**, 147–155, <https://doi.org/10.1007/s10735-017-9713-8>
- 7 Bielefeld, K.A., Amini-Nik, S. and Alman, B.A. (2013) Cutaneous wound healing: recruiting developmental pathways for regeneration. *Cell. Mol. Life Sci.* **70**, 2059–2081, <https://doi.org/10.1007/s00018-012-1152-9>
- 8 Assis-Ribas, T., Forni, M.F., Winnischofer, S.M.B., Sogayar, M.C. and Trombetta-Lima, M. (2018) Extracellular matrix dynamics during mesenchymal stem cells differentiation. *Dev. Biol.* **437**, 63–74, <https://doi.org/10.1016/j.ydbio.2018.03.002>
- 9 Kariminekoo, S., Movassaghpour, A., Rahimzadeh, A., Talebi, M., Shamsasenjan, K. and Akbarzadeh, A. (2016) Implications of mesenchymal stem cells in regenerative medicine. *Artif Cells Nanomed. Biotechnol.* **44**, 749–757, <https://doi.org/10.3109/21691401.2015.1129620>
- 10 Ko, S.H., Nauta, A.C., Morrison, S.D., Hu, M.S., Zimmermann, A.S., Chung, M.T. et al. (2018) PHD-2 suppression in mesenchymal stromal cells enhances wound healing. *Plast. Reconstr. Surg.* **141**, 55e–67e, <https://doi.org/10.1097/PRS.0000000000003959>
- 11 Motegi, S.I. and Ishikawa, O. (2017) Mesenchymal stem cells: the roles and functions in cutaneous wound healing and tumor growth. *J. Dermatol. Sci.* **86**, 83–89, <https://doi.org/10.1016/j.jdermsci.2016.11.005>
- 12 Han, Y., Jia, L., Zheng, Y. and Li, W. (2018) Salivary exosomes: emerging roles in systemic disease. *Int. J. Biol. Sci.* **14**, 633–643, <https://doi.org/10.7150/ijbs.25018>
- 13 Liu, X., Wang, S., Wu, S., Hao, Q., Li, Y., Guo, Z. et al. (2018) Exosomes secreted by adipose-derived mesenchymal stem cells regulate type I collagen metabolism in fibroblasts from women with stress urinary incontinence. *Stem Cell Res. Ther.* **9**, 159, <https://doi.org/10.1186/s13287-018-0899-9>
- 14 Toh, W.S., Lai, R.C., Hui, J.H.P. and Lim, S.K. (2017) MSC exosome as a cell-free MSC therapy for cartilage regeneration: implications for osteoarthritis treatment. *Semin. Cell Dev. Biol.* **67**, 56–64, <https://doi.org/10.1016/j.semcdb.2016.11.008>
- 15 Li, X., Xie, X., Lian, W., Shi, R., Han, S., Zhang, H. et al. (2018) Exosomes from adipose-derived stem cells overexpressing Nrf2 accelerate cutaneous wound healing by promoting vascularization in a diabetic foot ulcer rat model. *Exp. Mol. Med.* **50**, 29, <https://doi.org/10.1038/s12276-018-0058-5>
- 16 Dalirfardouei, R., Jamialahmadi, K., Jafarian, A.H. and Mahdipour, E. (2019) Promising effects of exosomes isolated from menstrual blood-derived mesenchymal stem cell on wound-healing process in diabetic mouse model. *J. Tissue Eng Regen. Med.* **13**, 555–568, <https://doi.org/10.1002/term.2799>
- 17 Ma, T., Fu, B., Yang, X., Xiao, Y. and Pan, M. (2019) Adipose mesenchymal stem cell-derived exosomes promote cell proliferation, migration, and inhibit cell apoptosis via Wnt/beta-catenin signaling in cutaneous wound healing. *J. Cell Biochem* **120**, 10847–10854, <https://doi.org/10.1002/jcb.28376>
- 18 Song, J., Su, Z.Z. and Shen, Q.M. (2020) Long non-coding RNA MALAT1 regulates proliferation, apoptosis, migration and invasion via miR-374b-5p/SRSF7 axis in non-small cell lung cancer. *Eur. Rev. Med. Pharmacol. Sci.* **24**, 1853–1862, <https://doi.org/10.26355/eurev/202002/20363>
- 19 Bhat, O.M., Yuan, X., Camus, S., Salloum, F.N. and Li, P.L. (2020) Abnormal lysosomal positioning and small extracellular vesicle secretion in arterial stiffening and calcification of mice lacking mucopolipin 1 gene. *Int. J. Mol. Sci.* **21**, 1713, <https://doi.org/10.3390/ijms21051713>
- 20 Santoso, M.R., Ikeda, G., Tada, Y., Jung, J.H., Vaskova, E., Sierra, R.G. et al. (2020) Exosomes from induced pluripotent stem cell-derived cardiomyocytes promote autophagy for myocardial repair. *J. Am. Heart Assoc.* **9**, e014345, <https://doi.org/10.1161/JAHA.119.014345>
- 21 Zhao, B., Zhang, Y., Han, S., Zhang, W., Zhou, Q., Guan, H. et al. (2017) Exosomes derived from human amniotic epithelial cells accelerate wound healing and inhibit scar formation. *J. Mol. Histol.* **48**, 121–132, <https://doi.org/10.1007/s10735-017-9711-x>
- 22 Shen, T., Zheng, Q., Luo, H., Li, X., Chen, Z., Song, Z. et al. (2020) Exosomal miR-19a from adipose-derived stem cells suppresses differentiation of corneal keratocytes into myofibroblasts. *Aging* **12**, 4093–4110, <https://doi.org/10.18632/aging.102802>
- 23 Ren, L., Chunxia, W., Kui, L. and Zuneng, L. (2018) LncRNA MALAT1 up-regulates VEGF-A and ANGPT2 to promote angiogenesis in brain microvascular endothelial cells against oxygen-glucose deprivation via targeting miR-145. *Biosci. Rep.* **2019**, BSR20180226, <https://doi.org/10.1042/BSR20180226>
- 24 Guo, S.C., Tao, S.C., Yin, W.J., Qi, X., Yuan, T. and Zhang, C.Q. (2017) Exosomes derived from platelet-rich plasma promote the re-epithelization of chronic cutaneous wounds via activation of YAP in a diabetic rat model. *Theranostics* **7**, 81–96, <https://doi.org/10.7150/thno.16803>
- 25 Cheon, S.S., Wei, Q., Gurung, A., Youn, A., Bright, T., Poon, R. et al. (2006) Beta-catenin regulates wound size and mediates the effect of TGF-beta in cutaneous healing. *FASEB J.* **20**, 692–701, <https://doi.org/10.1096/fj.05-4759com>
- 26 Jun, J.I. and Lau, L.F. (2018) Resolution of organ fibrosis. *J. Clin. Invest.* **128**, 97, <https://doi.org/10.1172/JCI93563>
- 27 Yu, J., Wang, M.Y., Tai, H.C. and Cheng, N.C. (2018) Cell sheet composed of adipose-derived stem cells demonstrates enhanced skin wound healing with reduced scar formation. *Acta Biomater.* **77**, 191–200, <https://doi.org/10.1016/j.actbio.2018.07.022>
- 28 Dimarino, A.M., Caplan, A.I. and Bonfield, T.L. (2013) Mesenchymal stem cells in tissue repair. *Front. Immunol.* **4**, 201, <https://doi.org/10.3389/fimmu.2013.00201>

- 29 Lin, Y.C., Grahovac, T., Oh, S.J., Ieraci, M., Rubin, J.P. and Marra, K.G. (2013) Evaluation of a multi-layer adipose-derived stem cell sheet in a full-thickness wound healing model. *Acta Biomater.* **9**, 5243–5250, <https://doi.org/10.1016/j.actbio.2012.09.028>
- 30 Nauta, A.J. and Fibbe, W.E. (2007) Immunomodulatory properties of mesenchymal stromal cells. *Blood* **110**, 3499–3506, <https://doi.org/10.1182/blood-2007-02-069716>
- 31 Kim, W.S., Park, B.S., Sung, J.H., Yang, J.M., Park, S.B., Kwak, S.J. et al. (2007) Wound healing effect of adipose-derived stem cells: a critical role of secretory factors on human dermal fibroblasts. *J. Dermatol. Sci.* **48**, 15–24, <https://doi.org/10.1016/j.jdermsci.2007.05.018>
- 32 Zhang, L., Xu, P., Wang, X., Zhang, M., Yan, Y., Chen, Y. et al. (2017) Activin B regulates adipose-derived mesenchymal stem cells to promote skin wound healing via activation of the MAPK signaling pathway. *Int. J. Biochem. Cell Biol.* **87**, 69–76, <https://doi.org/10.1016/j.biocel.2017.04.004>
- 33 Ong, H.T., Redmond, S.L., Marano, R.J., Atlas, M.D., von Unge, M., Aabel, P. et al. (2017) Paracrine activity from adipose-derived stem cells on in vitro wound healing in human tympanic membrane keratinocytes. *Stem Cells Dev.* **26**, 405–418, <https://doi.org/10.1089/scd.2016.0204>
- 34 Hu, L., Wang, J., Zhou, X., Xiong, Z., Zhao, J., Yu, R. et al. (2016) Exosomes derived from human adipose mesenchymal stem cells accelerates cutaneous wound healing via optimizing the characteristics of fibroblasts. *Sci. Rep.* **6**, 32993, <https://doi.org/10.1038/srep32993>
- 35 Ji, P., Diederichs, S., Wang, W., Boing, S., Metzger, R., Schneider, P.M. et al. (2003) MALAT-1, a novel noncoding RNA, and thymosin beta4 predict metastasis and survival in early-stage non-small cell lung cancer. *Oncogene* **22**, 8031–8041, <https://doi.org/10.1038/sj.onc.1206928>
- 36 Schmidt, L.H., Spieker, T., Koschmieder, S., Schaffers, S., Humberg, J., Jungen, D. et al. (2011) The long noncoding MALAT-1 RNA indicates a poor prognosis in non-small cell lung cancer and induces migration and tumor growth. *J. Thorac. Oncol.* **6**, 1984–1992, <https://doi.org/10.1097/JTO.0b013e3182307eac>
- 37 Tano, K., Mizuno, R., Okada, T., Rakwal, R., Shibato, J., Masuo, Y. et al. (2010) MALAT-1 enhances cell motility of lung adenocarcinoma cells by influencing the expression of motility-related genes. *FEBS Lett.* **584**, 4575–4580, <https://doi.org/10.1016/j.febslet.2010.10.008>
- 38 Liu, J.Y., Yao, J., Li, X.M., Song, Y.C., Wang, X.Q., Li, Y.J. et al. (2014) Pathogenic role of lncRNA-MALAT1 in endothelial cell dysfunction in diabetes mellitus. *Cell Death Dis.* **5**, e1506, <https://doi.org/10.1038/cddis.2014.466>
- 39 Sun, L., Sun, P., Zhou, Q.Y., Gao, X. and Han, Q. (2016) Long noncoding RNA MALAT1 promotes uveal melanoma cell growth and invasion by silencing of miR-140. *Am. J. Transl. Res.* **8**, 3939–3946
- 40 Tang, Y., Jin, X., Xiang, Y., Chen, Y., Shen, C.X., Zhang, Y.C. et al. (2015) The lncRNA MALAT1 protects the endothelium against ox-LDL-induced dysfunction via upregulating the expression of the miR-22-3p target genes CXCR2 and AKT. *FEBS Lett.* **589**, 3189–3196, <https://doi.org/10.1016/j.febslet.2015.08.046>
- 41 Zhang, G., Song, K. and Yan, H. (2019) MicroRNA-124 represses wound healing by targeting SERP1 and inhibiting the Wnt/beta-catenin pathway. *Adv. Clin. Exp. Med.* **28**, 711–718, <https://doi.org/10.17219/acem/94163>
- 42 Yang, P., Bu, P. and Li, C. (2017) miR-124 inhibits proliferation, migration and invasion of malignant melanoma cells via targeting versican. *Exp. Ther. Med.* **14**, 3555–3562, <https://doi.org/10.3892/etm.2017.4998>
- 43 Yang, H.L., Tsai, Y.C., Korivi, M., Chang, C.T. and Hseu, Y.C. (2017) Lucidone promotes the cutaneous wound healing process via activation of the PI3K/AKT, Wnt/beta-catenin and NF-kappaB signaling pathways. *Biochim. Biophys. Acta Mol. Cell Res.* **1864**, 151–168, <https://doi.org/10.1016/j.bbamcr.2016.10.021>
- 44 Lu, T., Bao, Z., Wang, Y., Yang, L., Lu, B., Yan, K. et al. (2016) Karyopherinbeta1 regulates proliferation of human glioma cells via Wnt/beta-catenin pathway. *Biochem. Biophys. Res. Commun.* **478**, 1189–1197, <https://doi.org/10.1016/j.bbrc.2016.08.093>





ADSC



Supplementary Figure S1 Spindle ADSCs were observed through the microscope.
×40 magnification.






Supplementary Figure S2 Transmission electron microscopy photomicrographs of exosomes. Scale bar, 100 nm.



Supplementary Figure S3 The Caspase3, Bax and Bcl-2 expression were detected by western blot assay.




Supplementary Figure S4 Cell proliferation of HaCaT and HDF cells was evaluated by CCK-8 assay.






Supplementary Figure S5 The capacity of cell migration in HaCaT and HDF cells was analyzed by the scratch wound healing assay.




Hacat






HDF



Supplementary Figure S6 Apoptosis of HaCaT and HDF cells treated with H₂O₂, H₂O₂+ADSC-Exo, H₂O₂+ADSC-Exo-shNC or H₂O₂+ADSC-Exo-shMALAT1 were monitored by flow cytometry assay.



Supplementary Figure S7 Migration of HaCaT and HDF cells treated with H₂O₂, H₂O₂+ADSC-Exo, H₂O₂+ADSC-Exo-shNC or H₂O₂+ADSC-Exo-shMALAT1 were analyzed by the scratch wound healing assay.



Supplementary Figure S8 Migration of HaCaT and HDF cells treated with FH535

was performed using the scratch wound healing assay.