

Research Article

# Retroviral insertional mutagenesis implicates E3 ubiquitin ligase RNF168 in the control of cell proliferation and survival

Aytug Kizilers<sup>1</sup>, Mark R. Pickard<sup>2,3</sup>, Cathleen E. Schulte<sup>1</sup>, Kiren Yacqub-Usman<sup>2,4</sup>, Nicola J. McCarthy<sup>2,5</sup>, Shu-Uin Gan<sup>1,6</sup>, David Darling<sup>1</sup>, Joop Gäken<sup>1</sup>, Gwyn T. Williams<sup>2</sup> and Farzin Farzaneh<sup>1</sup>

<sup>1</sup>Department of Haematological Medicine, King's College London, The Rayne Institute, London SE5 9NU, U.K.; <sup>2</sup>Apoptosis Research Group, School of Life Sciences, Huxley Building, Keele University, Keele ST5 5BG, U.K.; <sup>3</sup>Institute of Medicine, Bache Hall, University of Chester, Chester CH2 1BR, U.K.; <sup>4</sup>Tumour and Vasculature Biology, School of Medicine, University of Nottingham, Queen's Medical Centre, Nottingham NG7 2UH, U.K.; <sup>5</sup>Horizon Discovery Ltd, 7100 Cambridge Research Park, Beach Road, Waterbeach, Cambridge CB25 9TL, U.K.; <sup>6</sup>Department of Surgery, National University of Singapore, Singapore 117597

**Correspondence:** Gwyn T. Williams (g.t.williams@keele.ac.uk)



The E3 ubiquitin ligase RNF168 is a ring finger protein that has been previously identified to play an important regulatory role in the repair of double-strand DNA breaks. In the present study, an unbiased forward genetics functional screen in mouse granulocyte/macrophage progenitor cell line FDCP1 has identified E3 ubiquitin ligase RNF168 as a key regulator of cell survival and proliferation. Our data indicate that RNF168 is an important component of the mechanisms controlling cell fate, not only in human and mouse haematopoietic growth factor dependent cells, but also in the human breast epithelial cell line MCF-7. These observations therefore suggest that RNF168 provides a connection to key pathways controlling cell fate, potentially through interaction with PML nuclear bodies and/or epigenetic control of gene expression. Our study is the first to demonstrate a critical role for RNF168 in the mechanisms regulating cell proliferation and survival, in addition to its well-established role in DNA repair.

## Introduction

The expression, replication and repair of DNA are pivotal to the physiological functioning of all cells. Understanding of the molecular mechanisms, control and integration of these processes is far from complete and this area is the subject of a great deal of research. This is justified, not only by the obvious importance of these processes in cellular physiology, but also by the many examples of abnormalities in these processes contributing to oncogenesis (reviewed by Hanahan and Weinberg [1]). The existence of important connections between DNA damage sensing mechanisms (DNA damage response (DDR)) and control of gene expression and their effect on processes such as cell proliferation and differentiation have been suspected for several decades [2]. There is now direct molecular evidence for some of these mechanisms. These range from DNA strand break mediated activation of p53 inhibiting haematopoietic progenitor cell differentiation [3,4], and the induction of differentiation by specific DDR genes such as *GADD45A* in haematopoietic stem cells [5], to the involvement of CDK12 in the regulation of DDR and embryonic development [6] as well as damage-induced modulation of miRNAs that affect cell cycle progression, apoptosis and differentiation [7-9].

Ongoing progress in our understanding of gene expression, DNA replication and repair most often relies on detailed investigation of previously identified molecules and, as a consequence, generally progresses incrementally. By contrast, forward genetics strategies allow unbiased approaches that can identify key molecules involved in rate-limiting steps independently through the subversion of individual gene function [10]. Successful forward genetics strategies include cDNA functional expression cloning [11-16]

Received: 23 May 2017  
Revised: 11 July 2017  
Accepted: 27 July 2017

Accepted Manuscript Online:  
27 July 2017  
Version of Record published:  
14 August 2017

and retroviral insertional mutagenesis (RIM) [16–20]. Indeed, current RIM studies have focused attention on the role of E3 ubiquitin ligase RNF168 in the control of cell fate.

Post-translational modification of proteins is extensively involved in controlling cell behaviour. Addition of ubiquitin to target proteins, either as a monomer or in the form of ubiquitin chains, is now recognized to have many important regulatory roles in addition to the targeting of proteins for degradation by the proteasome [21,22]. In particular, ubiquitination of nuclear proteins plays a central role both in DNA repair [22–24] and in epigenetic control of gene expression [25–27], including the expression of tumour suppressor genes [27].

Extensive studies have implicated RNF168 in the repair of double-strand DNA breaks [23,28–32]. The repair of double-strand DNA breaks is a complex process in which RNF168 and RNF8 catalyse the ubiquitination of histone H2A subtypes that leads to recruitment of protein components of the DNA repair machinery, including 53BP1 and BRCA1 [28–32]. Mutation in RNF168 produces RIDDLE syndrome in humans [33], although some of the features of the phenotype, such as craniofacial abnormalities and short stature, have hitherto been difficult to ascribe to aberrant DNA repair alone.

Although *RNF168* is amplified in some cancers [32,34], the observations reported below are the first to demonstrate the involvement of this gene in the control of cell survival and proliferation. Most recently, RNF168 has been shown to regulate PML nuclear bodies (PML NBs) [35], suggesting a potential mechanism for the regulation of proliferation and apoptosis by RNF168 described below.

## Materials and methods

### Materials

Recombinant mouse interleukin-3 (mIL-3) was obtained from R&D Systems (Abingdon, U.K.) and recombinant human interleukin-3 (hIL-3), reagents for real-time quantitative RT-PCR (RT-qPCR), Lipofectamine 2000 and the pcDNA3.1 and TopoPCR2.1 vectors were from Life Technologies Ltd (Paisley, U.K.). Cell culture reagents were from the latter source or from Sigma–Aldrich (Poole, U.K.). The plasmid pCMVSPORT6-RNF168 (MGC: 45398; IMAGE 5163887), which contains the complete coding sequence of human RNF168, was from Source BioScience (Nottingham, U.K.) and nucleofector solution T was from Lonza Bioscience (Verviers, Belgium). QuikChange<sup>®</sup> XL Site-directed Mutagenesis Kit was from Agilent Technologies (Stockport, U.K.) and polybrene was from Sigma–Aldrich (Poole, U.K.). siRNAs #1–#4 to human RNF168 (product codes: #1, Hs\_FLJ35794\_1; #2, Hs\_RNF168\_2; #3, Hs\_FLJ35794\_3; #4, Hs\_RNF168\_1) were from Qiagen Ltd (Crawley, U.K.); negative control (NC) siRNA (product 102728) and HiPerFect reagent were also from the latter source. The MTS assay kit (CellTiter 96 AQueous One Solution Cell Proliferation Assay) was from Promega (Southampton, U.K.) and the Muse Cell Cycle Assay Kit was from Millipore (U.K.) Ltd (Watford, U.K.). Protein Assay Kit II and precast gels were from Bio–Rad Laboratories (Hemel Hempstead, U.K.). The RNF168 and  $\beta$ -actin antibodies for immunoblotting were from Abcam (Cambridge, U.K.), whereas the anti-myc and FITC-labelled anti-mouse IgG antibodies for immunofluorescence were from Santa Cruz Biotechnology (Heidelberg, Germany) and Sigma–Aldrich (Poole, U.K.) respectively. Hybond-P PVDF membranes were from Amersham Biosciences (Little Chalfont, U.K.).

### Cell culture

The mouse haematopoietic granulocyte/macrophage progenitor cell line FDCP1 [36–38] was maintained in RPMI-1640 medium supplemented with 10% FBS, 2 mM L-glutamine, 100 U/ml penicillin, 100  $\mu$ g/ml streptomycin and 1 ng/ml recombinant mIL-3. Cells were deprived of mIL-3 by centrifugation and resuspension in mIL-3-free medium for two cycles of washing and cloning in soft agar without mIL-3. 293T cells were maintained in DMEM medium containing 10% FBS, 100 U/ml penicillin and 100  $\mu$ g/ml streptomycin. TF-1 cells were routinely maintained in R-10 medium (comprises RPMI-1640 containing 2 mM L-glutamine, 1 mM sodium pyruvate, 10 mM HEPES, 10% FBS and 50  $\mu$ g/ml gentamicin) supplemented with recombinant hIL-3 (5 ng/ml) and MCF7 [39] cells were maintained in R-10 medium; all cells were cultured at 37°C in a humidified incubator with 5% CO<sub>2</sub>.

### Generation of IL-3-independent FDCP-1B cell clones by RIM

In the present study, the P3MP packaging cell line which produces M3Pneo-sup virus was used for infection of FDCP1B cells. For RIM, a total of  $1 \times 10^8$  cells were co-cultured with the irradiated P3MP packaging cells for 2 days in the presence of mIL-3. Then the cells were harvested and incubated overnight in fresh medium also in the presence of interleukin-3 (IL-3). Next, while  $1 \times 10^8$  cells were seeded in soft agar in the absence of mIL-3,  $1 \times 10^8$  cells were taken through to the next round of co-culture with the P3MP packaging cell line. These cycles of co-culture were repeated for a total number of 15 times. After each cycle,  $1 \times 10^8$  cells were cloned in soft agar in the absence of

mIL-3. A total of 95 IL-3 independent FDCP1B clones were isolated at the end of 15 rounds of co-culture. The IL-3 independent FDCP1B clones were initially expanded in 1 ml culture in the absence of mIL-3 and G418. Following their successful expansion, G418 selection was performed on all clones, each of which was found to be G418 resistant. In the present study, we have analysed one of these clones, referred to as PAP60 [40].

## Inverse PCR

The inverse PCR (I-PCR) method was performed as described previously [41]. Briefly, genomic DNA of PAP60 cells was digested with the *Sau3A* restriction enzyme, followed by ligation and subsequent *XbaI* digestion. The sequence flanking the 3'-LTR of the integrated provirus in PAP60 was obtained using the outward directed primers MPSVA1 (AAACTGCTGAGGGCGGGACC) and MPSVA2 (AGTTCGCTTCTCGCTTCTGT). The PCR fragments produced by I-PCR were cloned into the TopoPCR2.1 vector and the genomic DNA at the 3' end of the integrated provirus flanking sequence was sequenced, i.e. 5'GTAATTTCTTCTTTTAGCACTAAGAAGCTTTAGAAAAGCTTTGTTAGGAGAAGGGTAGCCTAAAGAATACTGAAAGAATAATTACAAAAATTCCTGTTCGGATC3' and identified by a BLAST search of the human genome.

## Plasmids and transfection

### Transduction into FDCP1B

All pLKO.1-shRNA plasmids were designed by The RNAi Consortium with the following clone reference numbers: control shRNA targeting GFP (shRNA-eGFP (SHC005)) and shRNA targeting mouse RNF168; shRNF168A (TRCN0000040876), shRNF168B (TRCN0000040873) (Open BioSystems).

The coding sequence for full-length human RNF168 (IMAGE: BC046815) was subcloned into pcDNA3.1 plasmid in frame with a C-terminal Myc and His fusion domain; RNF168WT. The point mutations in the RING (RNF168H33A) and MIU domains (RNF168A179G/A450G) were introduced by site-directed mutagenesis using the QuikChange<sup>®</sup> XL Site-directed Mutagenesis Kit. All constructs were sequence verified.

### Transfection into TF-1 cells

TF-1 cells ( $2 \times 10^6$ ) in 0.1 ml nucleofector solution T, were nucleofected with 2  $\mu$ g plasmid constructs comprising pCMVSPORT6-RNF168 or empty pCMVSPORT6 vector (for controls) using program G-016, and cells were immediately plated in 3 ml I-20 medium (comprises Iscove's modified Dulbecco's medium containing 2 mM L-glutamine, 20% FBS and 50  $\mu$ g/ml gentamicin) supplemented with hIL-3 (5 ng/ml) and allowed to recover for 20 h. Cells were then washed twice with R-10 medium, and resuspended ( $3 \times 10^5$  cells/ml) in R-10 medium  $\pm$  hIL-3 (5 ng/ml) and cultured for a further 22 h before counting.

## RNAi

### shRNA (LVshRNA)

Lentiviral vector was produced by transient co-transfection of 293T cells with lentiviral packaging plasmids; pCMV $\Delta$ 8.91 and pMDG2 with the target lentiviral constructs; pLKO.1-shRNAs, using a calcium phosphate co-precipitation method. Supernatant, containing the vector, was harvested at 48 h (harvest I) and 72 h (harvest II). The vector supernatant was first clarified through a 0.45-  $\mu$ m filter and concentrated by centrifugation at 10000g (Beckman J2-MC) overnight at 4°C, resuspended in 500  $\mu$ l RPMI-1640 medium and stored at -80°C. Cells were infected with lentivirus vectors in the presence of 4  $\mu$ g/ml polybrene. A multiplicity of infection (MOI) of 3 was used for infection.

### siRNA

To determine the effect of RNF168 knockdown on culture growth, four siRNAs which target different portions of RNF168 sequence were individually studied, along with NC siRNA for controls. For TF1 cells, complexes were prepared by mixing 9  $\mu$ l HiPerFect reagent with 209  $\mu$ l siRNA (430 nM in Opti-MEM-I). After 15 min, these were added dropwise to cells ( $4 \times 10^5$  cells in 0.2 ml R-10 medium with 5 ng/ml IL-3; 12-well plates). After a further 3 h, R-10 medium with 5 ng/ml IL-3 (0.8 ml) was added, and cells were cultured for a further 70 h to allow silencing to occur. Cells were then reseeded ( $3 \times 10^5$  cells/ml in R-10 medium with 5 ng/ml hIL-3) and cultured for a further 72 h to assay the growth parameters. For MCF7 cells, a fast-forward transfection protocol was employed: complexes were prepared by mixing 12  $\mu$ l HiPerFect reagent with 100  $\mu$ l siRNA (120 nM in Opti-MEM-I). After 15 min, these were added dropwise to freshly trypsinized, exponentially growing cells ( $1.5 \times 10^5$  cells in 2.3 ml R-10 medium; six-well plates) and these were cultured for 70 h. Cells were then replated in either fresh R-10 medium for MTS assay (0.5

$\times 10^4$  cells/96-well plate), direct cell counting and/or cell cycle analysis (both at  $0.8 \times 10^5$  cells/12-well plate) at the indicated times, or in R-10 medium supplemented with 10% cell-conditioned medium (400 cells/six-well plate) for clonogenic assays.

## RT-qPCR

Total RNA was isolated using the TRIzol<sup>®</sup> reagent and 1  $\mu$ g of purified total RNA from each sample was reverse transcribed to cDNA with oligo-dT primers by using a SuperScript II<sup>™</sup> Kit according to the manufacturer's protocol. Quantitative PCRs were performed with Platinum<sup>®</sup> SYBR<sup>®</sup> Green qPCR SuperMix-UDG Kit (Thermo Fisher Scientific) using an Applied Biosystems 7900 HT thermal cycler. The following primers were used: mouse RNF168 (5'-AGGCAGGTCTGAGGAGAAAGTGTT-3', 5'-AGGCAGGTCTGAGGAGAAAGTGTT-3'), mouse Smc1 (5'-ACCATGGAAGTGAATATTGTATACAGCTAT-3', 5'-AACTGCTGGGTCAAAGGTAAC-3') and  $\beta$ -actin (5'-TGAACCCTAAGGCCAACCGTGA-3', 5'-AGTCCATCACAATGCCTGTGGTACG-3'). Cycling conditions were 5 min at 95°C followed by 40 cycles of 15 s at 95°C, 30 s at 55°C and 30 s at 72°C. The  $\Delta C_T$  method was applied to estimate relative transcript levels. Levels of  $\beta$ -actin amplification were used as an endogenous reference to normalize each sample.

## Growth analyses

Direct cell counting was performed after Nigrosin Blue (0.1% (w/v)) staining using a haemocytometer and light microscopy; cells which excluded the dye were considered to be viable. MTS assays were performed according to the manufacturer's instructions. For clonogenic assays, the colonies formed following 3 weeks of culture were counted after staining with Crystal Violet (0.5% (w/v) in methanol). Cell cycle analysis was performed on a Muse Cell Analyzer using a Muse Cell Cycle Assay Kit according to the supplied instructions.

## Immunofluorescence

293T cells were cultured on glass coverslips in six-well plates and transfected with pcDNA3.1/RNF168WT, RNF168H33A or RNF168A179G/A450G plasmid using Lipofectamine 2000 according to the manufacturer's instructions. Thirty-six hours after transfection, cells were fixed in 4% paraformaldehyde for 30 min and permeabilized in 0.2% Triton-X 100 solution for 15 min followed by blocking with 3% BSA/PBS for 30 min. For RNF168 protein detection by immunofluorescence, cells were incubated with mouse anti-myc primary antibody (1:200 dilution) for 1 h, after extensive washing, the cells were incubated with secondary antibody (FITC anti-mouse IgG, 1:100) for 1 h, followed by washing with PBS. Coverslips were air dried and counterstained with DAPI. Images were acquired by fluorescence microscopy.

## Western blot analysis

Whole protein lysates were extracted using RIPA lysis buffer supplemented with PMSF and protease inhibitor cocktail; the concentration of isolated proteins was determined using Protein Assay Kit II. Protein (50  $\mu$ g) was electrophoresed (10% precast gel), then transferred to Hybond-P PVDF membranes. These membranes were incubated with anti-RNF168 in 5% skimmed milk and appropriate secondary antibodies. Blots were then stripped and reprobed with an antibody to  $\beta$ -actin. Western blot imaging and quantification were carried out using the LI-COR ECL system (LI-COR, Lincoln, U.S.A.).


## Statistical analyses

Data are presented as the mean and S.E.M.; the number of observations ( $n$ ) refers to different transfected samples, each derived from a separate culture of cells. Data were analysed by either one-way ANOVA, using either Bonferroni's multiple comparison test (MCT) or Dunnett's MCT (the latter when comparing multiple groups compared with a single group) for post hoc analysis, or by two-way ANOVA, using Dunnett's MCT. Homogeneity of variance was checked by Bartlett's test and, where necessary, data were transformed (log or square root) prior to analysis. All analyses were performed using GraphPad Prism v4.03.

## Results

### RIM of mouse haematopoietic cells

The FDCP1 cell line is a growth factor dependent haematopoietic granulocyte/macrophage cell line originally isolated from DBA/2 mouse bone marrow [36]. FDCP1 cells were cloned using soft agar and limiting dilution in the presence




**Figure 1. RIM clone PAP60 is IL-3 independent for both survival and proliferation but is not insensitive to IL-3**

(a) FDCP-1B and PAP60 cells were washed three times with RPMI-1640 medium lacking mIL-3 and then seeded at  $1 \times 10^5$  cells per ml in the presence or absence of 1 ng/ml mIL-3. Cell proliferation was measured by MTT assay at the indicated time points over a 5-day period. (b) The presence of IL-3 still exerts a strong protective affect for FDCP-1B and PAP60 cells when apoptosis is induced by etoposide treatments. FDCP-1B and PAP60 cells were washed and seeded at  $2 \times 10^5$  cells in 2 ml culture medium, in the presence or absence of mIL-3 with 3.4- $\mu$ M etoposide (Eto). After 24 h, cells were harvested, stained with Propidium Iodide, and apoptosis was detected using flow cytometry. Apoptotic bodies appear in the sub-G<sub>1</sub> peak. Data are the mean  $\pm$  S.D. of three independent experiments.

of IL-3 and the subclone FDCP1B was selected for subsequent analyses because this subclone undergoes apoptosis more rapidly and synchronously on withdrawal of IL-3 [37] than parental FDCP1 cells [38].

FDCP1B cells were infected with retrovirus M3Pneo-sup. M3Pneo-sup is a myeloproliferative sarcoma virus (MPSV)-based retroviral vector in which all the viral genes have been removed (splice acceptor sequences and upstream sequences necessary for efficient splicing of Mo-MuLV are retained) and the selectable marker gene, neomycin phosphotransferase (*neoR*) has been inserted in order to select cells that carry an integrated provirus [42-44]. The retrovirally infected FDCP1B cells were plated in soft agar in the absence of IL-3 and IL-3-independent clones were isolated. One of these clones, PAP60 [40], was selected for further study and showed a complete IL-3 independence for both survival and growth (Figure 1a). In order to determine whether the mutant cells could respond to IL-3 at all,



**Figure 2. Identification of the provirus integration site in mouse chromosome 16 and expression of neighbouring genes**

(a and c) Schematic representation of mouse chromosome 16, *Smco1* and *RNF168*. In the PAP60 cells, the provirus is integrated 229 bases upstream of the 5' transcription start site of the *RNF168* and 2452 bp downstream of the 3' end of the *Smco1* gene. (b) Relative expression of *Smco1* and *RNF168* in FDCP-1B and PAP60 cells. PAP60 and FDCP1B cells were grown in the presence (1 ng/ml) of mIL-3 for 48 h. The cells were then washed and grown in the presence or absence of mIL-3 for 12 h. mRNA expression of *Smco1* and *RNF168* in parental FDCP-1B cells and PAP60 cells in the presence (1 ng/ml) or absence of mIL-3 were analysed by reverse transcription and real-time PCR with specific primers to *Smco1*, *RNF168* and  $\beta$ -actin. The transcript levels are expressed relative to  $\beta$ -actin mRNA levels in the parental FDCP-1B cells. RT-qPCR measurements were repeated three times with similar results.

they were challenged with topoisomerase II inhibitor etoposide in the presence and absence of IL-3 (Figure 1b). The presence of IL-3 produced substantial protection against etoposide-induced apoptosis of both the PAP60 mutant cell line and the parental cell line, indicating that the PAP60 cell line retained responsiveness to IL-3 even though it was no longer IL-3-dependent for survival or proliferation.

The key advantage of the RIM strategy is that it allows the site of insertion of the provirus to be determined by I-PCR [41,45]. This technique revealed that in the PAP60 cells, the provirus insertion site is in mouse chromosome 16, 229 bases upstream of the transcription start site of the gene encoding *RNF168* (Figure 2a), and 2452 bases downstream of *Smco1* (2310010M20Rik), a 200-amino acid single-pass membrane protein. The human homologue of *Smco1* is a 214-amino acid protein known as *C3orf43* [46,47]. Since the effects of proviral integration are not entirely predictable, the transcription levels of the two genes flanking the integration site were determined by RT-qPCR in PAP60 cells and the parental FDCP1B cells, both with and without IL-3 (Figure 2b). The level of expression of *Smco1* was the same in PAP60 and parental FDCP1B cells. By contrast, the level of expression of *RNF168* was doubled in PAP60

cells relative to parental FDCP1B cells. The expression of the two genes examined was unaffected by the presence of IL-3 in both PAP60 and parental FDCP1B cells (Figure 2b). Since the exact mechanism by which retroviral insertion increases transcription from the nearby *RNF168* gene is not clear, *RNF168* mRNA levels were manipulated in other ways (see below).

### Analysis of the role of RNF168 in mouse cells

Since the RT-qPCR analysis above suggested that RNF168 may be involved in the IL-3-independence observed in the PAP60 cells, the effect of down-regulation of endogenous RNF168 expression was examined. PAP60 and parental FDCP1B cells were transduced with shRNA vectors targeting RNF168. shRNAs targeting GFP were used as a control. RNF168-targeting shRNAs reduced *RNF168* mRNA levels by 60–70% in both PAP60 and FDCP1B cells (Figure 3a). This was accompanied by substantial reductions in culture cell density for both RNF168 shRNAs (Figure 3b), further implicating RNF168 in the control of proliferation in these murine growth factor dependent cells. This reduction in cell culture growth by both shRNF168 constructs was confirmed in a time-course experiment over 5 days (Figure 3c).

It is well established that RNF168 plays a key role in the repair of double-strand DNA breaks through the ubiquitination of nuclear proteins [23,28,33,48–50]. In order to examine the possibility that RNF168 might also be involved in modification of proteins in other cellular locations, Myc-tagged RNF168 was expressed in 293T cells and examined by fluorescence microscopy. RNF168 could only be detected within the nucleus, with a speckled appearance (Supplementary Figure S1). This nuclear distribution is consistent with the recently reported association of RNF168 with PML-NB [35].

### Analysis of the role of RNF168 in human haematopoietic cells


In order to test the hypothesis that RNF168 might play a role in the regulation of growth factor dependence in human as well as mouse cells, we analysed the effect of overexpression of RNF168 in the human growth factor dependent cell line TF-1 [15]. TF-1 is a human haematopoietic cell line and TF-1 cells are dependent on IL-3 or GM-CSF for survival and proliferation [15]. TF-1 cells were transfected with a RNF168 expression construct, in order to overexpress RNF168, or with vector only (Figure 4a). Transfected cells were incubated in the presence or absence of IL-3 for 22 h; both culture viability and viable cell number in control vector transfected TF-1 cells were significantly reduced in the absence of IL-3. By contrast, culture viability and viable cell number were unaffected by IL-3 withdrawal in the RNF168 overexpressing TF-1 cells (Figure 4b–d), as seen for PAP60 cells (Figure 1), implicating RNF168 in the control of cell fate in human as well as mouse cells. Down-regulation of RNF168 in IL-3-supplemented TF-1 cells (Figure 4e) produced reductions in both the total cell density and the viable cell density of cultures (Figure 4f–h), further indicating a role for endogenous RNF168 in the proliferation of these human haematopoietic cells.

### Analysis of the role of RNF168 in human breast cancer cells

Since both the cell lines, mouse FDCP1B and human TF-1, are growth factor dependent haematopoietic cell lines, we extended the study to investigate the role of endogenous RNF168 in the proliferation and survival of human breast cell line MCF-7. MCF-7 cell cultures were treated with siRNAs specific for RNF168, and a non-targeting NC siRNA. Western blotting confirmed partial depletion of RNF168 protein (Figure 5a). RNF168 siRNAs significantly reduced MTS culture growth measurements and both viable cell density and total cell density (Figure 5b–e). In addition, RNF168 siRNAs substantially reduced the colony-forming ability of MCF-7 cells cloned for 72 h after treatment (Figure 5f). Cell cycle analysis indicated that depletion of endogenous RNF168 increased accumulation of cells in the G<sub>0</sub>/G<sub>1</sub> phase, with corresponding decrease in cells in S-phase and G<sub>2</sub>/M (Figure 5g–i). These consistent observations indicate that the involvement of RNF168 in the control of human cell proliferation is not restricted to haematopoietic growth factor dependent cells.

## Discussion

A forward genetics strategy for the identification of functionally critical components of cell regulatory mechanisms has several important advantages: first, it is entirely independent of established knowledge; and second, it automatically focuses on elements that have controlling roles rather than on secondary phenomena [51–54]. RIM is an important application of the forward genetics strategy that has identified many unanticipated genes that encode key components of the mechanisms regulating cell fate. Many of these genes are implicated in oncogenesis and/or resistance to cancer therapies [18,19]. Our unbiased screen for genes implicated in growth factor independence in haematopoietic cells has identified an E3 ubiquitin ligase RNF168.




**Figure 3. Down-regulation of RNF168 in FDCP-1B and PAP60 cells leads to reduced numbers of cells**

(a,b) FDCP-1B and PAP60 cells were transduced with LV-shRNF168A, LV-shRNF168B and LV-shGFP (MOI =3). (a) RNA levels and (b) cell density were determined on day 4 after transduction. Data are expressed as mean  $\pm$  S.D., and statistical analyses of the data were performed using Student's *t* test. Values of  $P < 0.05$  were considered to be statistically significant. The data are representative of three independent experiments for FDCP-1B and two independent experiments for PAP60 cells. (a) RNF168 RNA levels are expressed relative to the levels in the untreated corresponding cell line (PAP60 absolute level is two fold higher than FDCP1 absolute level Figure 2b). (c) Reduction in cell number by down-regulation of RNF168. FDCP-1B and PAP60 cells were transduced with LV-shRNF168A, LV-shRNF168B and LV-shGFP vectors, at an MOI of 3. Cell numbers were determined by MTT assay over an 8-day period. Data are expressed as a percentage MTT activity compared with non-infected cells.


RNF168 has a well-established and important role in the modification of chromosomal proteins required for the repair of double-strand DNA breaks [23,28-32]. The observations presented above strongly suggest a broader role in the regulation of cell fate, i.e. in controlling cell survival and proliferation (Figures 1, 3-5). It is arguable that the amplification of *RNF168* in tumours is consistent with this wider role [32,34]. The observation that overexpression of





**Figure 4. RNF168 regulates the growth and survival of the human growth factor dependent cell line, TF1**

Cells were either transfected with a plasmid encoding human RNF168 or siRNA to human RNF168 and then cultured in the absence or presence of IL-3. (a–d) Cells were nucleofected with the plasmids pCMVSPORT6-RNF168 (R) or pCMVSPORT6 alone (V) and maintained in the presence of IL-3 for 20 h. (a) Western blot analysis demonstrating up-regulation of RNF168. Cells were then washed and replated at equal densities in medium containing (+) or without (–) IL-3 and cultured for a further 22 h before determining culture viability (b), viable cell number (c) and total cell number (d). Note that vector-transfected cells only demonstrate IL-3 dependence and that overexpression of RNF168 increases culture growth/viability in the absence of IL-3 only (\* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.001$  for intraplasmid comparisons; ^ $P < 0.05$  and ^^ $P < 0.001$  for interplasmid comparisons; one-way ANOVA and Bonferroni's MCT;  $n = 5$ ). (e–h) Cells were transfected with one of four different siRNAs to RNF168 (#1–#4) or with NC siRNA and maintained in the presence of IL-3 for 70 h. (e) Western blot analysis demonstrating down-regulation of RNF168. Cells were then washed and replated at equal densities in medium containing IL-3 and cultured for a further 70 h before determining culture viability (f), viable cell number (g) and total cell number (h). Note that silencing of RNF168 reduces culture growth (\* $P < 0.05$  and \*\* $P < 0.01$  compared with NC siRNA-transfected cells; one-way ANOVA and Dunnett's MCT;  $n = 5$ ).



**Figure 5. RNF168 also regulates the growth of human growth factor independent cells**

The breast cancer cell line, MCF7, was transfected with one of the four different siRNAs to RNF168 (#1–#4) or with NC siRNA and after 70 h, cells were replated at equal densities for analyses of culture growth and the cell cycle. (a) Western blot analysis demonstrating the effectiveness of the treatments. (b–d) Culture growth parameters determined by direct microscopic counting at 48 h *post*-replating. Note that silencing of RNF168 has negligible effect on culture viability (b) but significantly reduces viable (c) and total (d) cell numbers ( $*P < 0.05$  and  $**P < 0.01$  compared with NC siRNA-transfected cells; one-way ANOVA and Dunnett's MCT;  $n \geq 3$ ). (e) MTS assay of culture growth up to 96 h *post*-plating confirming reduced growth of cells transfected with siRNA to RNF168; statistical analysis (two-way ANOVA and Dunnett's MCT;  $n \geq 3$ ) demonstrates different ( $P < 0.05$ ) values for all siRNAs at 72 h and for siRNAs #3 and #4 at 96 h. (f) Clonogenic assay data demonstrating that long-term growth of cells is compromised following silencing of RNF168 ( $*P < 0.05$  and  $**P < 0.01$  compared with NC siRNA-transfected cells; one-way ANOVA and Dunnett's MCT;  $n \geq 3$ ). (g–i) Cell cycle analysis at 24 h *post*-replating, indicating reduced proliferation following silencing of RNF168 ( $*P < 0.05$ ,  $**P < 0.01$  and  $***P < 0.001$  compared with NC siRNA-transfected cells; one-way ANOVA and Dunnett's MCT;  $n = 3$ ).

RNF168 confers growth factor independence in both human and mouse haematopoietic cells (Figures 1, 3–5) suggests its involvement at key stages in pathways controlling cell fate. Moreover, our data suggest that this involvement of RNF168 in cell proliferation and survival is not limited to the cells of the haematopoietic lineage as reduction in the endogenous expression of RNF168 significantly affects the survival, proliferation and colony-forming ability of human breast epithelial cell line MCF-7.

Almost all previous reports of RNF168 activity have been concerned with the modification of nuclear proteins [31,55] and, consistent with this, our analysis of the subcellular localization of RNF168 (Supplementary Figure S1) indicates nuclear localization in patterns similar to those recently reported by Shire et al. [35]. In this context, it is important to appreciate that the post-translational modification of chromosomal proteins, including by ubiquitination [56,57], is involved in the regulation of many cellular processes, and this includes gene expression as well as DNA repair. For example, Bhatnagar et al. [27] reported that the ubiquitination of H2A by the E3 ubiquitin ligase TRIM37 resulted in the down-regulation of tumour suppressor genes in breast epithelial cells.

Several recent papers have shown that RNF168's role extends beyond histone modification [35,58,59]. As well as PML [35], RNF168-mediated ubiquitination of FOXM1 (transcription factor forkhead box M1 [58] and TOP2 $\alpha$  (topoisomerase II $\alpha$  [59]) has also been demonstrated. The observation that RNF168 expression affects the accumulation of proteins in PML NBs may prove to be especially significant [35]. Down-regulation of RNF168 increased the level of PML NBs and overexpression of RNF168 produced a corresponding decrease in PML NBs [35]. PML is a well-established tumour suppressor [60–64] and a reduction or loss in expression of PML in other systems, e.g. in PML<sup>-/-</sup> mice, results in increased proliferation and reduced apoptosis [60–63]. In particular, in TF1 cells, expression of the PML–RAR  $\alpha$  fusion protein, which acts as dominant-negative PML [64], protects these cells from apoptosis induced by growth factor withdrawal, allowing growth factor independent proliferation. Negative regulation of PML by RNF168 [35] could therefore account for the inhibition of apoptosis and growth factor independent proliferation that we report here. The functional importance of RNF168 in regulating cell proliferation and apoptosis may have important implications for the cell's response to DNA damage. Through connecting the DNA repair process to PML-NB activity, RNF168 may help to link the DDR to the subsequent fate of the cell.

In summary, an unbiased functional screen identified RNF168 in mouse haematopoietic cell lines that survived and proliferated in the absence of growth factor. Subsequent experiments suggest that RNF168 may play a key part of the mechanisms regulating cell survival and proliferation in both mouse and human cells, potentially through interaction with PML-NB.

## Competing interests

The authors declare that there are no competing interests associated with the manuscript.

## Author contribution

G.T.W. and F.F. conceived the study and drafted the manuscript with A.K. and M.R.P. A.K., C.E.S., S.-U.G., N.J.M., D.D. and J.G. carried out the experiments with mouse cell lines, and M.R.P. and K.Y.-U. carried out the experiments with human cell lines. All authors read and approved the final manuscript.

## Funding

This work was supported by the Bloodwise [grant number 08046]; the Breast Cancer Now [grants numbers 2011NovPR05 & 2013NovSP208]; the Rosetrees Trust; and the National Institute for Health Research (NIHR) Biomedical Research Centre (BRC) awarded to the Institute of Psychiatry in partnership with King's College Hospital NHS Trust and King's College London; and the Experimental Cancer Medicine Centre at King's College London.

## Abbreviations

BRCA1, Breast Cancer 1.; DDR, DNA damage response; hIL-3, human interleukin-3; IL-3, interleukin-3; I-PCR, inverse PCR; MCT, multiple comparison test; mIL-3, mouse interleukin-3; MOI, multiplicity of infection; NC, negative control; MoMuLV, Moloney Murine Leukemia Virus; PML, Promyelocytic Leukemia Protein; PML NB, PML nuclear body; RIM, retroviral insertional mutagenesis; RNF168, Ring Finger Protein 168.; RT-qPCR, real-time quantitative RT-PCR.

## References

- 1 Hanahan, D. and Weinberg, R.A. (2011) Hallmarks of cancer: the next generation. *Cell* **144**, 646–674
- 2 Farzaneh, F., Zalin, R., Brill, D. and Shall, S. (1982) DNA strand breaks and ADP-ribosyl transferase activation during cell-differentiation. *Nature* **300**, 362–366

- 3 Yamashita, M., Nitta, E. and Suda, T. (2016) Regulation of hematopoietic stem cell integrity through p53 and its related factors. *Ann. N.Y. Acad. Sci.* **1370**, 45–54
- 4 Mahfoudhi, E., Lordier, L., Marty, C., Pan, J., Roy, A., Roy, L. et al. (2016) DNA-damage response gene GADD45A induces differentiation in hematopoietic stem cells without inhibiting cell cycle or survival. *Oncotarget* **34**, 699–710
- 5 Wingert, S., Thalheimer, F.B., Haetscher, N., Rehage, M., Schroeder, T. and Rieger, M.A. (2016) DNA-damage response gene GADD45A induces differentiation in hematopoietic stem cells without inhibiting cell cycle or survival. *Stem Cells* **34**, 699–710
- 6 Juan, H.-C., Lin, Y., Chen, H.-R. and Fann, M.-J. (2016) Cdk12 is essential for embryonic development and the maintenance of genomic stability. *Cell Death Differ.* **23**, 1038–1048
- 7 Tutar, Y. (2014) miRNA and cancer; computational and experimental approaches. *Curr. Pharm. Biotechnol.* **15**, 429
- 8 Wright, C.M., Dan, T., Dicker, A.P. and Simone, N.L. (2014) microRNAs: the short link between cancer and RT-induced DNA damage response. *Front. Oncol.* **4**, 133
- 9 Shen, T., Sanchez, H.N., Zan, H. and Casali, P. (2015) Genome-wide analysis reveals selective modulation of microRNAs and mRNAs by histone deacetylase inhibitor in B cells induced to undergo class-switch DNA recombination and plasma cell differentiation. *Front. Immunol.* **6**, 627
- 10 Stark, G.R. and Gudkov, A.V. (1999) Forward genetics in mammalian cells: functional approaches to gene discovery. *Hum. Mol. Genet.* **8**, 1925–1938
- 11 Chambers, I., Colby, D., Robertson, M., Nichols, J., Lee, S., Tweedie, S. et al. (2003) Functional expression cloning of Nanog, a pluripotency sustaining factor in embryonic stem cells. *Cell* **113**, 643–655
- 12 Williams, G.T., Hughes, J.P., Stoneman, V., Anderson, C.L., McCarthy, N.J., Mourtada-Maarabouni, M. et al. (2006) Isolation of genes controlling apoptosis through their effects on cell survival. *Gene Ther. Mol. Biol.* **10**, 255–261
- 13 Mourtada-Maarabouni, M., Kirkham, L., Jenkins, B., Rayner, J., Gonda, T.J., Starr, R. et al. (2003) Functional expression cloning reveals proapoptotic role for protein phosphatase 4. *Cell Death Differ.* **10**, 1016–1024
- 14 Mourtada-Maarabouni, M., Hedge, V.L., Kirkham, L., Farzaneh, F. and Williams, G.T. (2008) Growth arrest in human T-cells is controlled by the non-coding RNA growth-arrest-specific transcript 5 (GAS5). *J. Cell Sci.* **121**, 939–946
- 15 Kitamura, T., Takaku, F. and Miyajima, A. (1991) IL-1 up-regulates the expression of cytokine receptors on a factor-dependent human hematopoietic-cell line, Tf-1. *Int. Immunol.* **3**, 571–577
- 16 Williams, G.T. and Farzaneh, F. (2004) The use of gene function to identify the rate-limiting steps controlling cell fate. *Cancer Immunol. Immunother.* **53**, 160–165
- 17 Haupt, Y., Alexander, W.S., Barri, G., Klinken, S.P. and Adams, J.M. (1991) Novel zinc finger gene implicated as Myc collaborator by retrovirally accelerated lymphomagenesis in E-Mu-Myc transgenic mice. *Cell* **65**, 753–763
- 18 Jonkers, J. and Berns, A. (1996) Retroviral insertional mutagenesis as a strategy to identify cancer genes. *Biochim. Biophys. Acta* **1287**, 29–57
- 19 Uren, A.G., Kool, J., Berns, A. and van Lohuizen, M. (2005) Retroviral insertional mutagenesis: past, present and future. *Oncogene* **24**, 7656–7672
- 20 Suzuki, T., Minehata, K., Akagi, K., Jenkins, N.A. and Copeland, N.G. (2006) Tumor suppressor gene identification using retroviral insertional mutagenesis in Blm-deficient mice. *EMBO J.* **25**, 3422–3431
- 21 Komander, D. and Rape, M. (2012) The ubiquitin code. *Annu. Rev. Biochem.* **81**, 203–229
- 22 Jackson, S.P. and Durocher, D. (2013) Regulation of DNA damage responses by ubiquitin and SUMO. *Mol. Cell* **49**, 795–807
- 23 Mattioli, F., Vissers, J.H.A., van Dijk, W.J., Ikpa, P., Citterio, E., Vermeulen, W. et al. (2012) RNF168 ubiquitinates K13-15 on H2A/H2AX to drive DNA damage signaling. *Cell* **150**, 1182–1195
- 24 Mattioli, F. and Sixma, T.K. (2014) Lysine-targeting specificity in ubiquitin and ubiquitin-like modification pathways. *Nat. Struct. Mol. Biol.* **21**, 308–316
- 25 Tavares, L., Dimitrova, E., Oxley, D., Webster, J., Poot, R., Demmers, J. et al. (2012) RYBP-PRC1 complexes mediate H2A ubiquitylation at polycomb target sites independently of PRC2 and H3K27me3. *Cell* **148**, 664–678
- 26 Ben-Saadon, R., Zaaroor, D., Ziv, T. and Ciechanover, A. (2006) The polycomb protein Ring1B generates self atypical mixed ubiquitin chains required for its *in vitro* histone H2A ligase activity. *Mol. Cell* **24**, 701–711
- 27 Bhatnagar, S., Gazin, C., Chamberlain, L., Ou, J., Zhu, X., Tushir, J.S. et al. (2014) TRIM37 is a new histone H2A ubiquitin ligase and breast cancer oncoprotein. *Nature* **516**, 116–120
- 28 Gatti, M., Pinato, S., Maiolica, A., Rocchio, F., Prato, M.G., Aebersold, R. et al. (2015) RNF168 promotes noncanonical K27 ubiquitination to signal DNA damage. *Cell Rep.* **10**, 226–238
- 29 Smeenk, G., Wiegant, W.W., Marteiijn, J.A., Luijsterburg, M.S., Sroczynski, N., Costelloe, T. et al. (2013) Poly(ADP-ribosylation) links the chromatin remodeler SMARCA5/SNF2H to RNF168-dependent DNA damage signaling. *J. Cell Sci.* **126**, 889–903
- 30 Hu, Y., Wang, C., Huang, K., Xia, F., Parvin, J.D. and Mondal, N. (2014) Regulation of 53BP1 protein stability by RNF8 and RNF168 is important for efficient DNA double-strand break repair. *PLoS ONE* **9**, e110522
- 31 Munoz, M.C., Laulier, C., Gunn, A., Cheng, A., Robbiani, D.F., Nussenzweig, A. et al. (2012) RING finger nuclear factor RNF168 is important for defects in homologous recombination caused by loss of the breast cancer susceptibility factor BRCA1. *J. Biol. Chem.* **287**, 40618–40628
- 32 Zong, D., Callen, E., Pegoraro, G., Lukas, C., Lukas, J. and Nussenzweig, A. (2015) Ectopic expression of RNF168 and 53BP1 increases mutagenic but not physiological non-homologous end joining. *Nucleic Acids Res.* **43**, 4950–4961
- 33 Stewart, G.S., Panier, S., Townsend, K., Al-Hakim, A.K., Kolas, N.K., Miller, E.S. et al. (2009) The RIDDLE syndrome protein mediates a ubiquitin-dependent signaling cascade at sites of DNA damage. *Cell* **136**, 420–434
- 34 <http://cancersangeracuk/cosmic/gene/analysis?In=RNF168> (2016) Wellcome Trust Sanger Institute
- 35 Shire, K., Wong, A.I., Tatham, M.H., Anderson, O.F., Ripsman, D., Gulstene, S. et al. (2016) Identification of RNF168 as a PML nuclear body regulator. *J. Cell Sci.* **129**, 580–591

- 36 Dexter, T.M., Garland, J., Scott, D., Scolnick, E. and Metcalf, D. (1980) Growth of factor-dependent hematopoietic precursor cell-lines. *J. Exp. Med.* **152**, 1036–1047
- 37 McCarthy, N.J. (1993) Apoptosis Induced by Cancer Chemotherapeutic Drugs and its Genetic Suppression.. *Ph.D. Thesis*, University of Birmingham
- 38 Williams, G.T., Smith, C.A., Spooncer, E., Dexter, T.M. and Taylor, D.R. (1990) Hematopoietic colony stimulating factors promote cell-survival by suppressing apoptosis. *Nature* **343**, 76–79
- 39 Soule, H.D., Vazquez, J., Long, A., Albert, S. and Brennan, M. (1973) Human cell line from a pleural effusion derived from a breast carcinoma. *J. Natl. Cancer Inst.* **51**, 1409–1416
- 40 Schulte, C.E. (1996) Isolation and Analysis of Factor-independent Murine Haematopoietic Progenitor Cells (FDCP-1) by Retroviral Insertional Mutagenesis. *Ph.D. Thesis*, University of London
- 41 Sambrook, J. and Russell, D.W. (2006) Inverse PCR. *Cold Spring Harb. Protoc.* **2006**, doi:10.1101/pdb.prot3487
- 42 Kollek, R., Stocking, C., Smadajoffe, F. and Ostertag, W. (1984) Molecular-cloning and characterization of a leukemia-inducing myeloproliferative sarcoma-virus and 2 of its temperature-sensitive mutants. *J. Virol.* **50**, 717–724
- 43 Laker, C., Stocking, C., Bergholz, U., Hess, N., Delamarter, J.F. and Ostertag, W. (1987) Autocrine stimulation after transfer of the granulocyte macrophage colony-stimulating factor gene and autonomous growth are distinct but interdependent steps in the oncogenic pathway. *Proc. Natl. Acad. Sci. U.S.A.* **84**, 8458–8462
- 44 Stocking, C., Bergholz, U., Friel, J., Klingler, K., Wagener, T., Starke, C. et al. (1993) Distinct classes of factor-independent mutants can be isolated after retroviral mutagenesis of a human myeloid stem-cell line. *Growth Factors* **8**, 197–209
- 45 Nakamura, T. (2005) Retroviral insertional mutagenesis identifies oncogene cooperation. *Cancer Sci.* **96**, 7–12
- 46 Skarnes, W.C., Rosen, B., West, A.P., Koutsourakis, M., Bushell, W., Iyer, V. et al. (2011) A conditional knockout resource for the genome-wide study of mouse gene function. *Nature* **474**, 337–342
- 47 Gerhard, D.S., Wagner, L., Feingold, E.A., Shenmen, C.M., Grouse, L.H., Schuler, G. et al. (2004) The status, quality, expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). *Genome Res.* **14**, 2121–2127
- 48 Doil, C., Mailand, N., Bekker-Jensen, S., Menard, P., Larsen, D.H., Pepperkok, R. et al. (2009) RNF168 binds and amplifies ubiquitin conjugates on damaged chromosomes to allow accumulation of repair proteins. *Cell* **136**, 435–446
- 49 Pinato, S., Gatti, M., Scanduzzi, C., Confalonieri, S. and Penengo, L. (2011) UMI, a novel RNF168 ubiquitin binding domain involved in the DNA damage signaling pathway. *Mol. Cell Biol.* **31**, 118–126
- 50 Gudjonsson, T., Altmeyer, M., Savic, V., Toledo, L., Dinant, C., Grofte, M. et al. (2012) TRIP12 and UBR5 suppress spreading of chromatin ubiquitylation at damaged chromosomes. *Cell* **150**, 697–709
- 51 Kile, B.T. and Hilton, D.J. (2005) The art and design of genetic screens: mouse. *Nat. Rev. Genet.* **6**, 557–567
- 52 Kool, J. and Berns, A. (2009) High-throughput insertional mutagenesis screens in mice to identify oncogenic networks. *Nat. Rev. Cancer* **9**, 389–399
- 53 Touw, I.P. and Erkeland, S.J. (2007) Retroviral insertion mutagenesis in mice as a comparative oncogenomics tool to identify disease genes in human leukemia. *Mol. Ther.* **15**, 13–19
- 54 Ranzani, M., Annunziato, S., Adams, D.J. and Montini, E. (2013) Cancer gene discovery: exploiting insertional mutagenesis. *Mol. Cancer Res.* **11**, 1141–1158
- 55 Fradet-Turcotte, A., Canny, M.D., Escribano-Diaz, C., Orthwein, A., Leung, C.C.Y., Huang, H. et al. (2013) 53BP1 is a reader of the DNA-damage-induced H2A Lys 15 ubiquitin mark. *Nature* **499**, 50–54
- 56 Shanbhag, N.M., Rafalska-Metcalf, I.U., Balane-Bolivar, C., Janicki, S.M. and Greenberg, R.A. (2010) ATM-dependent chromatin changes silence transcription in cis to DNA double-strand breaks. *Cell* **141**, 970–981
- 57 Lukas, J., Lukas, C. and Bartek, J. (2011) More than just a focus: the chromatin response to DNA damage and its role in genome integrity maintenance. *Nat. Cell Biol.* **13**, 1161–1169
- 58 Kongsema, M., Zona, S., Karunarathna, U., Cabrera, E., Man, E.P.S., Yao, S. et al. (2016) RNF168 cooperates with RNF8 to mediate FOXM1 ubiquitination and degradation in breast cancer epirubicin treatment. *Oncogenesis* **5**, e252
- 59 Guturi, K.K.N., Bohgaki, M., Bohgaki, T., Srikumar, T., Ng, D., Kumareswaran, R. et al. (2016) RNF168 and USP10 regulate topoisomerase II alpha function via opposing effects on its ubiquitylation. *Nat. Commun.* **7**, 12638
- 60 Wang, Z.G., Ruggero, D., Ronchetti, S., Zhong, S., Gaboli, M., Rivi, R. et al. (1998) PML is essential for multiple apoptotic pathways. *Nat. Genet.* **20**, 266–272
- 61 Trotman, L.C., Alimonti, A., Scaglioni, P.P., Koutcher, J.A., Cordon-Cardo, C. and Pandolfi, P.P. (2006) Identification of a tumour suppressor network opposing nuclear Akt function. *Nature* **441**, 523–527
- 62 Salomoni, P., Ferguson, B.J., Wyllie, A.H. and Rich, T. (2008) New insights into the robe of PML in tumour suppression. *Cell Res.* **18**, 620–638
- 63 Tang, M.K., Liang, Y.J., Chan, J.Y.H., Wong, S.W., Chen, E., Yao, Y. et al. (2013) Promyelocytic leukemia (PML) protein plays important roles in regulating cell adhesion, morphology, proliferation and migration. *PLoS ONE* **8**, e59477
- 64 Fu, S.Q., Consoli, U., Hanania, E.G., Zu, Z.F., Claxton, D.F., Andreoff, M. et al. (1995) PML/RAR alpha, a fusion protein in acute promyelocytic leukemia, prevents growth factor withdrawal-induced apoptosis in TF-1 cells. *Clin. Cancer Res.* **1**, 583–590

## Supplemental Material For Publication

**Supplementary Figure 1.** Subcellular localization of RNF168. Myc-tagged wild-type (RNF168<sup>WT</sup>), RING domain mutated (RNF168<sup>H33A</sup>) and MIU domains mutated (RNF168<sup>A179G/A450G</sup>) RNF168 plasmids were transfected into 293T cells. RNF168 protein was visualized using anti-myc antibody; nuclei were counter stained with DAPI. RNF168<sup>WT</sup> (A) and RNF168<sup>H33A</sup> (B) proteins localized to nucleus with a dot-like, speckled pattern, while RNF168<sup>A179G/A450G</sup> (C) distributed evenly throughout the nucleus.

