

## Review Article

# Pivotal neuroinflammatory and therapeutic role of high mobility group box 1 in ischemic stroke

Seidu A. Richard<sup>1,2</sup>, Marian Sackey<sup>3</sup>, Zhaoliang Su<sup>1</sup> and Huaxi Xu<sup>1</sup>

<sup>1</sup>Department of Immunology, Jiangsu University, 301 Xuefu Road, Zhenjiang 212013, P.R. China; <sup>2</sup>Department of Surgery, Volta Regional Hospital, P.O. Box MA-374, Ho, Ghana-West Africa; <sup>3</sup>Department of Pharmacy, Volta Regional Hospital, P.O. Box MA-374, Ho, Ghana-West Africa

**Correspondence:** Seidu A. Richard (gbepoo@gmail.com)



Stroke is a major cause of mortality and disability worldwide. Stroke is a frequent and severe neurovascular disorder. The main cause of stroke is atherosclerosis, and the most common risk factor for atherosclerosis is hypertension. Therefore, prevention and treatment of stroke are crucial issues in humans. High mobility group box 1 (HMGB1) is non-histone nuclear protein that is currently one of the crucial proinflammatory alarmins in ischemic stroke (IS). It is instantly released from necrotic cells in the ischemic core and activates an early inflammatory response. HMGB1 may signal via its putative receptors, such as receptor for advanced glycation end products (RAGE), toll-like receptors (TLRs) as well as matrix metalloproteinase (MMP) enzymes during IS. These receptors are expressed in brain cells. Additionally, brain-released HMGB1 can be redox modified in the circulation and activate peripheral immune cells. The role of HMGB1 may be more complex. HMGB1 possesses beneficial actions, such as endothelial activation, enhancement of neurite outgrowth, and neuronal survival. HMGB1 may also provide a novel link for brain-immune communication leading to post-stroke immunomodulation. Therefore, HMGB1 is new promising therapeutic intervention aimed at promoting neurovascular repair and remodeling after stroke. In this review, we look at the mechanisms of secretion of HMGB1, the role of receptors, MMP enzymes, hypoglycemia, atherosclerosis, edema, angiogenesis as well as neuroimmunological reactions and post-ischemic brain recovery in IS. We also outline therapeutic roles of HMGB1 in IS.

## Introduction

Ischemic stroke (IS) occurs when a cerebral blood vessel is occluded. This disrupts the cerebral supply with nutrients and oxygen leading to permanent necrotic brain tissue destruction in the affected area of the respective terminal artery. This focal brain damage can result in various acute neurological deficits [1,2]. It is estimated that the overall costs of stroke care will account for 6.2% of the total burden of illness by 2020 [2,3]. Hypertension (HT), hyperlipidemia (HL), and diabetes mellitus (DM) are the most common risk factors for atherosclerosis, which is the key trigger of stroke [2,4].

High mobility group box 1 (HMGB1) is non-histone nuclear protein that is currently one of the crucial proinflammatory alarmins in IS [1]. It has been indicated that the secretion of endogenous alarmins also known as danger-associated molecular patterns (DAMPs) from necrotic cerebral cells may be one of the core initiators of the inflammatory cascade. It is now clear that DAMPs released from the necrotic tissues initiate local inflammatory response in the infarct core in ischemic brain injury but when released into the blood, it may signal peripheral immune cells [1,5]. Several studies have confirmed HMGB1 as one of the DAMPs that has been shown to be involved in ischemic brain injury [6-9]. It is usually confined in cell nuclei in the normal brain, translocates into the cytosol, and is secreted into the extracellular compartment under ischemic conditions. It has been reported that intracellular HMGB1 plays a critical role in activation and regulation of basal transcriptional machinery [8,10,11] and energy homeostasis. Studies have shown that the extracellular secretion of HMGB1 is observed within 2-4 h after

Received: 31 July 2017  
Revised: 29 September 2017  
Accepted: 18 October 2017

Accepted Manuscript Online:  
20 October 2017  
Version of Record published:  
5 December 2017

ischemia–reperfusion, increases vascular permeability, and promotes blood–brain barrier (BBB) breakdown [9,12]. Zhang et al. [12] indicated that the neutralization of extracellular HMGB1 prevented the BBB permeability and curtails the infarct volume in mice and rats. Therefore, HMGB1 is a fundamental DAMP in the hyperacute phase of ischemic brain injury. Qiu et al. [13] also indicated that the extracellular release of HMGB1 dwindles within 12 h after ischemia–reperfusion.

In this review, we look at the role of the redox form of HMGB1, the mechanisms of release of HMGB1 in the early phase of IS as well as the receptors via which HMGB1 interacts. We also looked at the role of HMGB1 in atherosclerosis, angiogenesis, edema, hyperglycemia as well as its neuroimmunological properties and post-ischemic brain recovery. We suggest that more research with both animal and human models is needed to further confirm that HMGB1 has therapeutic potential since initial studies proved that it promotes neurovascular repair and remodeling after stroke.

## The redox state of HMGB1 in strokes

HMGB1 is a non-histone DNA-binding protein of 215 amino acid residues arranged in three domains that include A box and B box domains arranged in an L-shape and a 30-amino acid long C-terminal tail [14–17]. Muhammad et al. [18] indicated that HMGB1 reduces ischemic brain injury in a mouse model of experimental stroke through Box A which they discovered as antagonist of inflammation during IS. Li et al revealed that the cytokine-inducing activity of disulphide HMGB1 is located on B the B-box and it is made up of Cys<sup>106</sup> residues. However, antibody-mediated targeting role of B box is still not known, so we suggest further focus on this role.

It has been shown that the redox modification of HMGB1 occurs at three cysteine residues at positions 23, 45, and 106, which regulate the functional properties of HMGB1. Increased levels of reactive oxygen species (ROS) lead to the formation of disulphide bonds between cysteine 23 and 45 [1]. These bonds have proven to induce nuclear factor  $\kappa$ -light-chain-enhancer of activated B cells (NF- $\kappa$ B) translocation to the nucleus and tumor necrosis factor alpha (TNF- $\alpha$ ) synthesis by macrophages [1,20]. Studies have further shown that all cysteine residues in HMGB1 can also be oxidized by the sustained ROS and nitrogen species (RNS) secretion from the mitochondria of apoptotic cells [1]. This kind of terminally oxidized HMGB1 is known as sulphonyl HMGB1, which gives anti-inflammatory signals and curtails excessive inflammatory activity [1,21]. HMGB1 can be modified into a reduced or oxidized form exerting specific cellular functions [1].

## HMGB1 release in the acute phase after stroke in animal models

Cerebral blood vessel occlusion in stroke causes a rapid necrosis of the affected brain region. Brain resident microglia and astrocytes in reaction to ischemic injury, get stimulated and release a storm of ROS and RNS and proinflammatory cytokines that cause secondary damage to the infarct area [1,22]. The onsite recruitment of peripheral immune cells that further exaggerate the oxidative stress through release of ROS and RNS is accompanied by extensive inflammation of the ischemic brain tissue. The inflammatory processes build a highly oxidative circumstance in the ischemic brain. Essentially, the action and function of several proteins is controlled by oxidation of cysteine, tyrosine, and methionine residues [1].

HMGB1 is expressed as a soluble peptide with proinflammatory cytokine function in reaction to a cell activation stimulus or passively after necrotic cell death. It synergizes with other proinflammatory molecules and cascades in the beginning of neuroinflammation [1]. Necrotic cells secrete hypoacetylated HMGB1 and signal innate immune cells to respond to the tissue injury that leads to the recruitment of mononuclear cells to the tissue injury and stimulate an inflammatory milieu [1,23]. HMGB1 immediately (i.e. within 1 h after ischemia onset) translocates from nuclei to the cytoplasm in neurones in the ischemic brain. Its release in activated microglia, astrocytes, and blood vessels is amplified 2 days after middle cerebral artery occlusion (MCAO) in rats [13,24]. *In vivo* knockdown of HMGB1 release by intrastriatal injection of HMGB1-shRNA 24 h before MCAO decreases infarct size and microglia activation, signifying a neurotoxic and proinflammatory role of HMGB1 after stroke [1,6].

HMGB1 expressed from necrotic tissue can induce inflammatory cytokine release in the infarcted brain. It has been demonstrated that stereotactical injection of fully reduced recombinant HMGB1 into the parietal cortex of mice forces the mRNA to release inducible nitric oxide synthase (iNOS) and interleukin (IL)-1 $\beta$  (IL-1 $\beta$ ) after 24 h [1,25]. Also *in vitro* cultures of mouse brain derived glial cells indicated amplified release of iNOS, IL-1 $\beta$ , Cyclooxygenase-2 (COX-2), and TNF- $\alpha$  after treatment with recombinant HMGB1 [25], signifying an activation of glial cells by HMGB1 that was potentially secreted by necrotic neurones. It is also clear that the ‘maturation’ of the HMGB1 in brain parenchyma and blood toward a hyperacetylated state can be detected within 24 h after brain insult [26].

## HMGB1 release in the acute phase after stroke in human models or patients

Studies have indicated that HMGB1 levels in the blood of stroke patients are higher as compared with matched healthy controls [27,28]. HMGB1 concentrations remain significantly elevated for a prolonged time period in symptomatic patients. Schulze et al. [29] demonstrated that in acute IS patients presented with elevated plasma concentrations of HMGB1 that persistently rise for 30 days after the primary insult and associated with a number of circulating leukocytes. They further indicated that increased HMGB1 levels have incoherent correlation with infarct size and location, since higher HMGB1 levels are observed in patients with more severe cerebral lesions [29].

Furthermore, the levels of HMGB1 in patients with stroke remain significantly elevated than in control subjects up to 14 days after the ischemic event while levels of the natural inhibitors of HMGB1, soluble receptor for advanced glycation end products (sRAGE) and endogenous secretory RAGE (esRAGE), remain indistinguishable from control subjects within 48 h following stroke. In patients with stroke, HMGB1 levels are adequately associated with IL-6 levels but not with the magnitude of brain tissue damage as evaluated by CT morphometry. Also, patients with stroke present an elevated fraction of activated positive cluster of differentiation 4 (CD4<sup>+</sup>) T cells in peripheral blood expressing interleukin-2 receptor alpha chain (CD25) or human leukocyte antigen-antigen D related (HLA-DR) when compared with controls [1,17,28]. Vogelgesang et al. [28] proposed the theory that HMGB1 acts as a link between brain tissue destruction by ischemic injury and the activation and type 1 T-helper (Th1) priming of T cells. They think it could be due to the similarity between the kinetics of serum HMGB1 and the kinetics observed for the absolute number of CD4<sup>+</sup> T cells expressing HLA-DR [1,17,28]. Studies have further demonstrated that higher serum concentrations of HMGB1 are associated with the severity of neurologic impairment and with elevated serum levels of TNF- $\alpha$  and IL-1 $\beta$  [1,17,26,30].

## Cerebral HMGB1 receptors in IS

HMGB1 modulates numerous neuroinflammatory events via ligation to different cell surface receptors. The main receptors for HMGB1 investigated in brain injury are the receptor for advanced glycation end products (RAGE) and toll-like receptors (TLR-2 and TLR-4) as well as matrix metalloproteinases (MMPs). These receptors are ubiquitously expressed on central nervous system (CNS) resident microglia, astrocytes, and neurones [31,32].


## RAGE

There is a growing body of evidence that the RAGE and its ligands are involved in the pathogenesis of various disorders, including cardiovascular, neurodegenerative, inflammatory, and autoimmune disorders [33]. In the brain, RAGE is expressed on glial cells and neurones [34-36]. Biopsy samples of human patients with unilateral cerebral infarction have shown elevated levels of RAGE [37]. In murine experimental stroke, RAGE levels are elevated in the ischemic brain hemisphere a day after the stroke [37]. Remarkably, serum levels of sRAGE are elevated in stroke patients in the acute phase [38].

A hypertensive challenge has an early and sustained effect on RAGE up-regulation in brain vessels of the cortex and hippocampus in mice [2,39] and plasma sRAGE levels are decreased in patients with essential HT [2,40], which means that HMGB1 and RAGE contribute to the pathogenesis of HT, the key risk factor for stroke (Figure 1). Also, HMGB1 and RAGE contributed to the pathogenesis of HL (Figure 1). This was evidenced when cell culture experiments using monocyte cell culture (U937) showed the relocation of HMGB1 from the nucleus to the cytoplasm under HL stress [2,41]. Furthermore, HMGB1 and RAGE play a role in the progression of DM [2,42], which is one of the risk factors for stroke (Figure 1). A study has proven that hyperglycemia accelerates the release of HMGB1 and RAGE in human aortic endothelial cells [2,43]. Studies have further demonstrated that accumulation of advanced glycation end products (AGEs) in DM is associated with the pathogenesis of vascular and inflammatory cell complications that characterize DM [1,2,44]. Hu et al. [45] indicated that AGE-induced autophagy of vascular smooth muscle cells (VSMCs) contributes to the process of AGE-induced proliferation of rat VSMCs, which is related to atherosclerosis in DM [1,2]. Haraba et al. [41] observed an elevation in serum HMGB1 levels in golden Syrian hamsters with induced HL. They further observed that HMGB1 release and RAGE expression were elevated in cultures of U937 cells exposed to hyperlipidemic sera [1,41].

## TLRs in IS

TLRs are a category of receptors for a variety of molecular patterns originated from microorganisms but also identify tissue-released alarmins [1,46,47]. TLRs are a set of type 1 transmembrane proteins and have ten members in humans



**Figure 1.** HMGB1 release during ischemic events interacts with TLR-2 and TLR-4 that is expression on monocytes via the adaptor protein MyD88 and elevates serum of TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 levels, which leads to cerebral vessel occlusion Also, HMGB1 release and RAGE expression increase risk factors (HP, DM, HL), which contributes significantly to cerebral vessel occlusion during IS.

(TLR1–TLR10) and 11 members in mice (TLR1–TLR9 and TLR-11 and TLR-13) [1]. Architecturally, they consist of an extracellular domain, a transmembrane domain, and a cytoplasmic domain. The extracellular domain is involved in ligand recognition while the cytoplasmic domain is essential for signal transduction via association with the adaptor protein called myeloid differentiation primary response gene [88] (MyD88) (Figure 1) [1,48].


Brasier [49] has demonstrated that TLR-mediated signaling induces the secretion of proinflammatory cytokines and the proliferation of different cells. A study has shown that boosted expression of TLRs on CNS resident innate immune cells and subsequent release of toxic molecules are detrimental for neuronal survival and activation of TLRs has been linked to the progression of IS [1,50]. In stroke patients, increased TLR-2 and TLR-4 expression on monocytes is associated with higher serum levels of TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 (Figure 1) [1,51]. TLR-2- and TLR-4-knockout mice showed less brain damage and neuronal deficits after 3 days of cerebral ischemia induced by MCAO [1,52,53]. Recently, Balosso et al. have shown that disulphide HMGB1 can increase *N*-Methyl-D-aspartic acid (NMDA)-induced neuronal cell death *in vitro* through its interaction with TLR-4 receptors [1,54]. The activation of microglia via TLR-4 induces amplification of neuronal cell loss and axonal damage after cerebral ischemia (Figure 1) [1,55].

## HMGB1 and MMPs in IS

MMPs are proteases that can break down extracellular proteins, such as collagen, and are involved in extracellular matrix remodeling. These proteases are usually located in the cytosol in an inactivated state and are cleaved by plasmin or other MMPs to their active state [56,57]. Research has shown that MMP-9 has direct cytotoxic effects and can disrupt the BBB [58]. Studies have proven that the major source of MMP is microglia and they are released following ischemia, especially MMP-3 and MMP-9 [59,60]. Pfefferkorn and Rosenberg [61] demonstrated that acute MMP inhibition reduces infarct size, brain edema, and recombinant tissue plasminogen activator (t-PA) induced hemorrhage (Figure 2) in experimental stroke models. Further studies proved that mice deficient in MMP-9 or MMP-3 have reduced ischemic injury relative to wild-type controls [62,63]. However, persistent inhibition of MMPs after ischemia may have detrimental effects on functional recovery, because these proteases are important in neurovascular remodeling after stroke [64].

Sapojnikova et al. [65] indicated a strong correlation between MMP-9 secretion and HMGB1 levels, which was moreover linked with poor outcome in human IS. Qiu et al. demonstrated that HMGB1 up-regulated MMP-9 in cultured neurones and astrocytes, as well as in mouse brain (Figure 2) [66]. They stated that this phenomenon was mediated primarily through the TLR4, which was highly expressed in neuronal and astrocyte cultures as well as mouse brain (Figure 2). Furthermore, HMGB1-induced MMP-9 responses were mostly suppressed after blockade of TLR4 signaling. Therefore, HMGB1 constitutively expressed on TLR-4 receptors in adjacent brain, thus up-regulating





**Figure 2. HMGB1 up-regulated MMP-9, which is mediated primarily through the TLR-4**

HMGB1 induced cytokines such as TNF- $\alpha$  and IL-1 $\beta$ , which may promote MMP-9 up-regulation indirectly. Up-regulation of MMPs leads to increased infarct size, brain edema, and recombinant t-PA induced hemorrhage, which accelerates neurovascular substrate damage including the tight junction protein Occludin. Similarly, anti-HMGB1 mAb suppresses the activity of MMP-9, with an inhibition of the increased permeability of BBB, in ischemic brain area and reduces neuronal and glial apoptosis.

MMP-9 and expanding neurovascular damage in ischemic brain injury. HMGB1 can also induce many other cytokines such as TNF- $\alpha$  and IL-1 $\beta$ , which may promote MMP-9 up-regulation indirectly (Figure 2) [66]. A potential source of elevated plasma MMP-9 and HMGB1 levels could be secretion from neutrophils, as MMP-9 is a neutrophil-derived blood MMP and a high quantity of HMGB1 characterizes these blood cells [65]. Activated MMP-9 could directly damage neurovascular substrates including the tight junction protein Occludin (Figure 2) [67]. Several studies have demonstrated that MMP-mediated degradation of Occludin is involved in the ischemia-induced BBB disruption [67-69]. Meanwhile therapeutically injected [70-72] and endogenous t-PA [73] have been implicated in the activation of MMP-9 during IS; HMGB1 may enhance the disruption of BBB structure through activation of MMPs via t-PA. Moreover, Liu et al. [7] indicated that anti-HMGB1 mAb suppresses the activity of MMP-9, with an inhibition of the increased permeability of BBB, in ischemic brain area (Figure 2). Inhibition of MMP-2 and MMP-9 reduces neuronal and glial apoptosis (Figure 2) [74].

## Atherosclerosis and HMGB1 in IS

Kalinina et al. [75] demonstrated that HMGB1 is essentially secreted by endothelial cells, smooth muscle cells and in CD68-positive macrophages situated close to the intima as well as in microvessels within the adventitia of human aorta. They further stated that active secretion of HMGB1 has been detected in areas adjacent to the necrotic core of atherosclerotic lesions [1,75]. Studies have shown that HMGB1 may be secreted from several cell types in the atherosclerotic plaque including smooth muscle cells, endothelial cells, foam cells, macrophages, and activated platelets [1,75-77]. HMGB1 when secreted, induces several inflammatory effects on endothelial cells, smooth muscle cells, and macrophages [17].

Studies have demonstrated that recombinant HMGB1 activates vascular endothelial cells initiating the release of intercellular adhesion molecule 1 (ICAM-1), vascular cell adhesion molecule 1 (VCAM-1), E-selectin, granulocyte colony stimulating factor (G-CSF), RAGE, TNF- $\alpha$ , monocyte chemotactic protein 1 (MCP-1), IL-8, plasminogen activator inhibitor 1, and t-PA [1,17,78-80]. Further studies have indicated that HMGB1 augments the formation of atherosclerotic plaques on smooth muscle cells as well as their proliferation, migration to the intimal layer, their release of more HMGB1 as well as C-reactive protein, and their secretion of MMP-2, MMP-3, and MMP-9 during IS [17,76,81]. The role of HMGB1 in the formation of atherosclerosis has been reported in apolipoprotein E deficient mice fed with a high-fat diet. Kanellakis et al. [82] reported that anti-HMGB1 neutralizing antibodies led to a dwindling of macrophage, dendritic cells (DC), and CD4<sup>+</sup> T-cell accumulation in atherosclerotic lesions and decreased the secretion of VCAM-1 and MCP-1 [17,80]. They indicated that neutralization of monoclonal antibodies against HMGB1 lessens atherosclerosis by 55%. HMGB1 also plays a key role in the pathogenesis of thrombosis. Studies

have shown that HMGB1 inhibits the anticoagulant protein C pathway via the thrombin–thrombomodulin complex, and stimulates release of tissue factor in monocytes *in vitro* [83,84]. Studies have further demonstrated that HMGB1 promotes development of microvascular thrombosis in rats [83,84].

## HMGB1 and angiogenesis in IS

The formation of new blood vessels in the late period of cerebral ischemia triggers a highly coupled neurorestorative process that mediates neurogenesis, synaptogenesis, and angiogenesis [1,85,86]. HMGB1 signaling can stimulate endothelial revival [79] and growth [87]. Researchers have shown that HMGB1 stimulates enlargement of endothelial progenitors, which boost angiogenesis; this outcome can be clogged by injection of HMGB1 siRNA 5 days post stroke [1,88,89]. HMGB1-mediated effect on angiogenesis may present a possible target for augmenting recovery several days after stroke onset. We therefore suggest that further studies are needed in this direction. Additionally, HMGB1 has been proven to accelerate neurite outgrowth and cell survival in neurones [1,90–92]. It has further been demonstrated that HMGB1 secreted by reactive astrocytes provides a signal for endothelial progenitor cell (EPC) mediated neurovascular remodeling, and this signaling between glial and vascular compartments may be critically significant for brain repair and stroke recovery [1,88]. Also, HMGB1 mediates cross-talk between reactive astrocytes and EPCs to promote functional recovery following focal cerebral ischemia [88].

## HMGB1 and edema formation in IS


Even though there are quite a lot of aquaporins (water channel proteins) expressed in the CNS, Aquaporin 4 (AQP4) is the major one in the brain, and is expressed in astroglial end feet around the surface of capillaries related to the BBB, glia limitans, and ependyma [93,94]. IS induces both immediate and delayed cell death, which is accompanied by a robust inflammatory response and cerebral edema that can exacerbate injury during reperfusion [94,95]. Brain edema is the main cause of death in patients with large infarction. AQP4, a water channel protein, is predominantly located in astroglial end feet [93,94] and has been shown to play an important role in water transport in the brain [94,96]. Also, AQP4 is considered to contribute to brain edema, particularly as AQP4-deficient mice have improved outcome following focal ischemic injury [94,97]. AQP4 may also contribute to cell adhesion [98] and migration [99], as well as neuroinflammation [94,100]. Interacting with other proinflammatory cytokines, extracellular HMGB1 aggravates inflammation and worsens injury in the ischemic brain, which further results in the up-regulation of aquaporins and exacerbation of brain edema. It is now clear that inhibition of HMGB1 is found to be neuroprotective for IS in rats [7,94]. We therefore propose that inhibition of HMGB1 could lead to reduction in AQP4 and hence reduction in brain edema.

## HMGB1 and neuroimmunological reactions in IS

Microglial cells are the resident macrophages of the brain and play an essential role as resident immunocomponent and phagocytic cells. It has been indicated that microglia activated under inflammation can transform into phagocytes and secrete a variety of mediators that are cytotoxic or cytoprotective [1,8]. Researchers have demonstrated that activated microglia after ischemia have the potential of secreting several proinflammatory cytokines such as TNF- $\alpha$ , IL-1, and IL-6, and nitric oxide ROS [1,101]. It has been proven that HMGB1 has essential roles for microglial activation minutes after cerebral ischemia [6] and may still be detected in microglia up to days after cerebral ischemia [24]. Although the infiltration of immune cells and the production of inflammatory mediators are conspicuous after IS, the direct activation of infiltrating immune cells is not achieved by HMGB1, which means that it may not be an ideal therapeutic target in the delayed phase of IS. Other researchers have proven that intravenous injection of neutralizing anti-HMGB1 mAb or intranasal delivery of HMGB1 siRNA provided a robust neuroprotection in the post-ischemic brain by antagonizing the proinflammatory function of HMGB1 [1,7,102].

## HMGB1 and hyperglycemia in IS

Studies have shown that glutamate, an excitatory amino acid, plays a central role in neuronal death [103,104]. Li et al. [105] demonstrated that hyperglycemia boosts extracellular glutamate accumulation in rats' models of forebrain ischemia [104]. Qiu et al. [13] discovered that glutamate induces the secretion of HMGB1 from neurones *in vitro* [104]. Huang et al. therefore proposed that hyperglycemia could worsen brain ischemic damage by boosting the early extracellular secretion of HMGB1 [104]. They indicated that hyperglycemia augments the early secretion of HMGB1 from ischemic brain tissue, which led to increased infarct volume, neurological deficit, cerebral edema, and BBB disruption [104]. Initial research finding has shown that BBB was significantly disrupted 4 h after reperfusion



**Figure 3. HMGB1 stimulates the production of IL-1, TNF- $\alpha$ , IL-6, and IL-8 and induces iNOS expression during ischemic brain damage**

Induction of iNOS and TNF- $\alpha$  occurs mainly in microglia. Induction of any of the factors above produces inflammatory response and the disruption of the BBB, leading to the aggravation of brain infarction.

in hyperglycemic rats [1,104,106,107]. Liu indicated that extracellular HMGB1 is involved in the disruption of BBB during the early phase of IS [7,104]. Disruption of BBB is a crucial event in the development of brain edema during the early stage of ischemic brain damage. Studies have shown that the development of brain edema is one of the most unfavorable ramifications of stroke and is greatly boosted during hyperglycemic stroke [1,104,106,108].

## HMGB1 and post-ischemic brain recovery

Currently activated T lymphocytes are perceived as the key contributors to secondary tissue damage and facilitate the progression of stroke outcome. Shichita et al. [109] observed an important role of  $\gamma$ - $\delta$  T cells intervening in inflammatory brain damage after cerebral ischemia. Liesz et al. [110] demonstrated that inhibition of leukocyte-invasion to brain after cerebral ischemia decreases neuroinflammation and ischemic tissue injury. Studies have also shown that the regulatory T- and B-cell populations are neuroprotective after brain ischemia [1,111-113]. Further studies have indicated that IL-10 has been successfully used as a therapeutic mediator to reduce post-stroke secondary neuroinflammation [1,112,114-116]. Moreover, higher plasma levels of IL-6 are taken as a predictive marker of poor outcome in stroke patients [117].

HMGB1 has been reported to stimulate the production of IL-1, TNF- $\alpha$ , IL-6, and IL-8 and to induce iNOS expression during ischemic brain damage (Figure 3) [1,118,119]. Further studies have shown that the expression of TNF- $\alpha$ , iNOS, and IL-1 $\beta$ , which were all up-regulated in the post-ischemic brain, was reduced when HMGB1 was inhibited [1,6,18]. Also, neuroprotective effect was accompanied by a reduction in the release of HMGB1 into the extracellular space and in the related inflammatory factors TNF- $\alpha$ , iNOS, IL-1 $\beta$ , and IL-6, both in the brain and serum of rats. The induction of iNOS and TNF- $\alpha$  following an ischemic insult was reported to occur mainly in microglia (Figure 3). Thus, it is likely that HMGB1 activates microglia in the brain, leading to the up-regulation of iNOS and TNF- $\alpha$  expression (Figure 3). The induction of iNOS and TNF- $\alpha$  has been reported to be involved in the inflammatory response and the disruption of the BBB, leading to the aggravation of brain infarction (Figure 3) [1,120,121]. The regulation of any one of these factors has been postulated to reduce ischemic injury.

Apoptosis is tightly regulated by the mitogen-activated protein kinase (MAPK) family, and the c-Jun N-terminal kinases (JNKs), extracellular signal-regulated kinase 1/2 (ERK1/2), and p38 members of this family have been demonstrated to be activated in ischemic reperfusion injury (I/R) [1,122]. MAPKs play important roles in transducing signals by phosphorylating intracellular enzymes, transcription factors, and cytosolic proteins involved in apoptosis and inflammatory cytokine production. Sustained MAPK activation has been shown to be associated with neuronal cell death/apoptosis following IS, and the inhibition of this pathway is neuroprotective [1,122].

## HMGB1 and therapeutic potentials of stroke medications

Therapeutic choices for acute stroke are still limited notwithstanding significant advances in the understanding of the pathophysiology of cerebral ischemia [123]. Statins have proven to attenuate the effects of HMGB1 on endothelial cells in two experimental studies. Yang et al. demonstrated that atorvastatin is able to inhibit endothelial activation *in vitro* upon HMGB1 stimulation in a dose-dependent manner [17,124]. They incubated endothelial cells with 10  $\mu$ M of atorvastatin and noticed that expression of ICAM-1 and E-selectin are reduced and inhibited HMGB1-stimulated leukocyte adhesion to endothelial cells. Furthermore, atorvastatin similarly suppressed HMGB1-induced TLR-4 expression and NF- $\kappa$ B nuclear translocation in endothelial cells [17,124]. Wang et al. further observed that atorvastatin had neuroprotective effect, and this effect is related to the decreased expression of HMGB1, its receptors (TLR-4 and RAGE), NF- $\kappa$ B in the acute phase of cerebral ischemia [17,125]. They therefore proposed that down-regulation of HMGB1-induced NF- $\kappa$ B activation pathway might be a potential mechanism of atorvastatin's neuroprotection in cerebral ischemia [17,125]. Haraba et al. [41] observed that fluvastatin reduced serum HMGB1 levels by 38.2% and led to a 1.46-fold reduction in HMGB1 mRNA expression in lung tissue [41]. The blockade of HMGB1 signaling with shRNA in the post-ischemic brain suppressed infarct size, microglial activation, and induction of proinflammatory mediators [6]. Parkkinen et al. [126] have demonstrated that HMGB1 has affinity for t-PA and accelerates its proteolytic activity.

Xiong et al. [94] have also demonstrated that administration of probenecid reduced infarct size, inhibited HMGB1 release from neurones, and down-regulated AQP4 expression in the cortical penumbra 48 h post stroke. They also observed that probenecid inhibits the activation of astrocytes and up-regulates AQP4 protein expression 48 h post stroke, which results in the attenuation of cerebral edema following focal ischemia. Because of its critical role in brain edema formation, AQP4 is regarded as an attractive therapeutic target for various brain disorders, including brain trauma, tumor, hydrocephalus, neuroinflammation, and focal cerebral ischemic injury [94,127]. Probenecid, as an inhibitor of Panx1 channels, inhibits ATP release induced by Panx1 activation [128,129], blocks inflammasome formation, and attenuates caspase 1 cleavage [130], thereby alleviating neuronal death and reducing brain inflammation and cerebral edema [94].

Administration of anti-HMGB1 neutralizing antibodies in experimental models of MCAO/reperfusion showed a significant decrease in infarct size and an improvement in neurologic deficits in treated rats. Anti-HMGB1 antibodies also prevented the increase in permeability of the BBB protecting the recipient from brain edema, inhibited activation of microglia and release of TNF- $\alpha$  and iNOS, while suppressing the activity of MMP-9 [1,7,12]. Additionally, the administration of numerous agents such as atorvastatin, minocycline, edaravone, cannabidiol, niaspan, and Tricin 7-glucoside was also shown to inhibit HMGB1 expression in brain ischemic tissue during the acute phase after stroke in experimental models of MCAO, so alleviating cerebral injury [1,125,131-135].

Tanshinone II A (Tan II A) effectively decreased neurologic impairment and tissue injury under cerebral ischemic conditions. This effect may be through down-regulation of HMGB1/HMGB1 receptors (TLR4 and RAGE)/NF- $\kappa$ B activation pathway, up-regulation of claudin-5 expression, and reduction in extravascular IgG. HMGB1 relation with NF- $\kappa$ B activation pathway may be one of Tan II A's effective therapeutic targets for cerebral ischemia. Tan II A inhibits HMGB1 translocation from nucleus to cytoplasm and/or extracellular space in the cortical peri-infarct regions [136].

Glycyrrhizin (GL) could be a new therapeutic possibility for the treatment of IS. Gong et al. [122] demonstrated that pretreatment with GL blocked and inhibited the extracellular cytokine activity of HMGB1 and explored the protective effect on I/R-induced apoptosis through the blockage of the JNK and p38-mediated pathways in rats *in vivo*. They stated that pretreatment with GL alleviated apoptosis injury resulting from cerebral I/R, and this was at least partly due to the inhibition of cytochrome *c* release and caspase 3 activity. Gwak et al. [137] further demonstrated that GL exhibited an anti-apoptotic effect by preventing HMGB1-induced cytochrome *c* release and caspase 3 activation *in vitro* in Huh-BAT cells (Korean Cell Line Bank, Seoul, Korea). Although the anti-apoptotic effect of GL has been linked with HMGB1 inhibition and caspase-dependent cytochrome *c* release, it is still not very clear how GL regulates caspase-dependent cytochrome *c* release through the inhibition of HMGB1. Some authors are of the view that GL



may modulate the activity of a particular kinase that contributes to cytochrome *c* translocation and is involved in I/R-induced HMGB1-dependent apoptosis [122].

## Conclusion

IS leads to the disruption of cerebral supply with nutrients and oxygen leading to permanent necrotic brain tissue destruction in the affected area of the respective terminal artery. HMGB1 presents dual effects after ischemic injury in the brain. HMGB1 acts as a proinflammatory mediator in the acute phase that amplifies damage in ischemic tissue through the activation of microglia, enhancement of inflammation, and increase in permeability of the BBB. In contrast, in the late phase after ischemic injury HMGB1 contributes to the recovery and remodeling process stimulating neurovascular repair mainly by astrocytes in the affected brain. We therefore suggest that more research with both animal and human models is needed to further confirm that HMGB1 has therapeutic potentials since initial studies proved that it promotes neurovascular repair and remodeling after stroke.

## Author contribution

S.A.R. is a guarantor of integrity of the entire concepts, design, and writing of the manuscript. M.S., Z.S., and H.X. contributed equally to the intellectual content of the manuscript.

## Competing interests

The authors declare that there are no competing interests associated with the manuscript.

## Funding

The authors declare that there are no sources of funding to be acknowledged.

## Abbreviations

AGE, advanced glycation end product; AQP4, aquaporin 4; BBB, blood–brain barrier; CNS, central nervous system; DAMP, danger-associated molecular pattern; DM, diabetes mellitus; EPC, endothelial progenitor cell; GL, glycyrrhizin; HL, hyperlipidemia; HLA-DR, human leukocyte antigen-antigen D related; HMGB1, high mobility group box 1; HT, hypertension; ICAM-1, intercellular adhesion molecule 1; IL-1 $\beta$ , interleukin-1 $\beta$ ; iNOS, inducible nitric oxide synthase; IS, ischemic stroke; JNK, c-Jun N-terminal kinase; MAPK, mitogen-activated protein kinase; MCAO, middle cerebral artery occlusion; MCP-1, monocyte chemotactic protein 1; MMP, matrix metalloproteinase; NF- $\kappa$ B, nuclear factor  $\kappa$ -light-chain-enhancer of activated B cell; RAGE, receptor for advanced glycation end product; RNS, reactive nitrogen species; ROS, reactive oxygen species; sRAGE, soluble RAGE; t-PA, tissue plasminogen activator; Tan II A, tanshinone II A; TLR, toll-like receptor; TNF- $\alpha$ , tumor necrosis factor  $\alpha$ ; VCAM-1, vascular cell adhesion molecule 1; VSMC, vascular smooth muscle cell.

## References

- 1 Singh, V., Roth, S., Veltkamp, R. and Liesz, A. (2016) HMGB1 as a key mediator of immune mechanisms in ischemic stroke. *Antioxid. Redox Signal.* **24**, 635–651
- 2 Kikuchi, K., Tancharoen, S., Ito, T., Morimoto-Yamashita, Y., Miura, N., Kawahara, K. et al. (2013) Potential of the angiotensin receptor blockers (ARBs) telmisartan, irbesartan, and candesartan for inhibiting the HMGB1/RAGE axis in prevention and acute treatment of stroke. *Int. J. Mol. Sci.* **14**, 18899–18924
- 3 Demaerschalk, B.M., Hwang, H.M. and Leung, G. (2010) US cost burden of ischemic stroke: a systematic literature review. *Am. J. Manag. Care* **16**, 525–533
- 4 Furie, K.L., Kasner, S.E., Adams, R.J., Albers, G.W., Bush, R.L., Fagan, S.C. et al. (2011) Guidelines for the prevention of stroke in patients with stroke or transient ischemic attack: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* **42**, 227–276
- 5 Richard, S.A., Xiang, L.-H., Yun, J.-X., Shanshan, Z., Jiang, Y.-Y., Wang, J. et al. (2017) Carcinogenic and therapeutic role of high-mobility group box 1 in cancer: is it a cancer facilitator, a cancer inhibitor or both? *World Cancer Res. J.* **4**, e919
- 6 Kim, J.B., Sig Choi, J., Yu, Y.M., Nam, K., Piao, C.S., Kim, S.W. et al. (2006) HMGB1, a novel cytokine-like mediator linking acute neuronal death and delayed neuroinflammation in the postischemic brain. *J. Neurosci.* **26**, 6413–6421
- 7 Liu, K., Mori, S., Takahashi, H.K., Tomono, Y., Wake, H., Kanke, T. et al. (2007) Anti-high mobility group box 1 monoclonal antibody ameliorates brain infarction induced by transient ischemia in rats. *FASEB J.* **21**, 3904–3916
- 8 Hayakawa, K., Qiu, J. and Lo, E.H. (2010) Biphasic actions of HMGB1 signaling in inflammation and recovery after stroke. *Ann. N. Y. Acad. Sci.* **1207**, 50–57
- 9 Shichita, T., Ago, T., Kamouchi, M., Kitazono, T., Yoshimura, A. and Ooboshi, H. (2012) Novel therapeutic strategies targeting innate immune responses and early inflammation after stroke. *J. Neurochem.* **123**, 29–38

- 10 Ellwood, K.B., Yen, Y.M., Johnson, R.C. and Carey, M. (2000) Mechanism for specificity by HMGB-1 in enhanceosome assembly. *Mol. Cell. Biol.* **20**, 4359–4370
- 11 Verrijdt, G., Haelens, A., Schoenmakers, E., Rombauts, W. and Claessens, F. (2002) Comparative analysis of the influence of the high-mobility group box 1 protein on DNA binding and transcriptional activation by the androgen, glucocorticoid, progesterone and mineralocorticoid receptors. *Biochem. J.* **361**, 97–103
- 12 Zhang, J., Takahashi, H.K., Liu, K., Wake, H., Liu, R., Maruo, T. et al. (2011) Anti-high mobility group box-1 monoclonal antibody protects the blood-brain barrier from ischemia-induced disruption in rats. *Stroke* **42**, 1420–1428
- 13 Qiu, J., Nishimura, M., Wang, Y., Sims, J.R., Qiu, S., Savitz, S.I. et al. (2008) Early release of HMGB-1 from neurons after the onset of brain ischemia. *J. Cereb. Blood Flow Metab.* **28**, 927–938
- 14 Read, C.M., Cary, P.D., Crane-Robinson, C., Driscoll, P.C. and Norman, D.G. (1993) Solution structure of a DNA-binding domain from HMGB1. *Nucleic Acids Res.* **21**, 3427–3436
- 15 Tsuda, K., Kikuchi, M., Mori, K., Waga, S. and Yoshida, M. (1988) Primary structure of non-histone protein HMGB1 revealed by the nucleotide sequence. *Biochemistry* **27**, 6159–6163
- 16 Seidu, R.A., Wu, M., Su, Z. and Xu, H. (2017) Paradoxical role of high mobility group box 1 in glioma: a suppressor or a promoter? *Oncol. Rev.* **11**, 325
- 17 de Souza, A.W., Westra, J., Limburg, P.C., Bijl, M. and Kallenberg, C.G. (2012) HMGB1 in vascular diseases: its role in vascular inflammation and atherosclerosis. *Autoimmun. Rev.* **11**, 909–917
- 18 Muhammad, S., Barakat, W., Stoyanov, S., Murikinati, S., Yang, H., Tracey, K.J. et al. (2008) The HMGB1 receptor RAGE mediates ischemic brain damage. *J. Neurosci.* **28**, 12023–12031
- 19 Li, J., Kokkola, R., Tabibzadeh, S., Yang, R., Ochani, M., Qiang, X. et al. (2003) Structural basis for the proinflammatory cytokine activity of high mobility group box 1. *Mol. Med.* **9**, 37–45
- 20 Yang, H., Lundback, P., Ottosson, L., Erlandsson-Harris, H., Venereau, E., Bianchi, M.E. et al. (2012) Redox modification of cysteine residues regulates the cytokine activity of high mobility group box-1 (HMGB1). *Mol. Med.* **18**, 250–259
- 21 Kazama, H., Ricci, J.E., Herndon, J.M., Hoppe, G., Green, D.R. and Ferguson, T.A. (2008) Induction of immunological tolerance by apoptotic cells requires caspase-dependent oxidation of high-mobility group box-1 protein. *Immunity* **29**, 21–32
- 22 Lipton, P. (1999) Ischemic cell death in brain neurons. *Physiol. Rev.* **79**, 1431–1568
- 23 Scaffidi, P., Misteli, T. and Bianchi, M.E. (2002) Release of chromatin protein HMGB1 by necrotic cells triggers inflammation. *Nature* **418**, 191–195
- 24 Kim, J.B., Lim, C.M., Yu, Y.M. and Lee, J.K. (2008) Induction and subcellular localization of high-mobility group box-1 (HMGB1) in the postischemic rat brain. *J. Neurosci. Res.* **86**, 1125–1131
- 25 Faraco, G., Fossati, S., Bianchi, M., Patrone, M., Pedrazzi, M., Sparatore, B. et al. (2007) High mobility group box 1 protein is released by neural cells upon different stresses and worsens ischemic neurodegeneration in vitro and in vivo. *J. Neurochem.* **103**, 590–603
- 26 Liesz, A., Dalpke, A., Mrcsko, E., Antoine, D.J., Roth, S., Zhou, W. et al. (2015) DAMP signaling is a key pathway inducing immune modulation after brain injury. *J. Neurosci.* **35**, 583–598
- 27 Goldstein, R.S., Gallowitsch-Puerta, M., Yang, L., Rosas-Ballina, M., Huston, J.M., Czura, C.J. et al. (2006) Elevated high-mobility group box 1 levels in patients with cerebral and myocardial ischemia. *Shock* **25**, 571–574
- 28 Vogelgesang, A., May, V.E., Grunwald, U., Bakkeboe, M., Langner, S., Wallaschofski, H. et al. (2010) Functional status of peripheral blood T-cells in ischemic stroke patients. *PLoS ONE* **5**, e8718
- 29 Schulze, J., Zierath, D., Tanzi, P., Cain, K., Shibata, D., Dressel, A. et al. (2013) Severe stroke induces long-lasting alterations of high-mobility group box 1. *Stroke* **44**, 246–248
- 30 Yang, Q.W., Lu, F.L., Zhou, Y., Wang, L., Zhong, Q., Lin, S. et al. (2011) HMGB1 mediates ischemia-reperfusion injury by TRIF-adaptor independent Toll-like receptor 4 signaling. *J. Cereb. Blood Flow Metab.* **31**, 593–605
- 31 Carty, M. and Bowie, A.G. (2011) Evaluating the role of Toll-like receptors in diseases of the central nervous system. *Biochem. Pharmacol.* **81**, 825–837
- 32 Ding, Q. and Keller, J.N. (2005) Evaluation of rage isoforms, ligands, and signaling in the brain. *Biochim. Biophys. Acta* **1746**, 18–27
- 33 Tang, S.C., Wang, Y.C., Li, Y.L., Lin, H.C., Manzanero, S., Hsieh, Y.H. et al. (2013) Functional role of soluble receptor for advanced glycation end products in stroke. *Arterioscler. Thromb. Vasc. Biol.* **33**, 585–594
- 34 Arancio, O., Zhang, H.P., Chen, X., Lin, C., Trinchese, F., Puzzo, D. et al. (2004) RAGE potentiates Abeta-induced perturbation of neuronal function in transgenic mice. *EMBO J.* **23**, 4096–4105
- 35 Bierhaus, A., Humpert, P.M., Morcos, M., Wendt, T., Chavakis, T., Arnold, B. et al. (2005) Understanding RAGE, the receptor for advanced glycation end products. *J. Mol. Med. (Berl.)* **83**, 876–886
- 36 Deane, R., Du Yan, S., Subramanian, R.K., LaRue, B., Jovanovic, S., Hogg, E. et al. (2003) RAGE mediates amyloid-beta peptide transport across the blood-brain barrier and accumulation in brain. *Nat. Med.* **9**, 907–913
- 37 Zhai, D.-X., Kong, Q.-F., Xu, W.-S., Bai, S.-S., Peng, H.-S., Zhao, K. et al. (2008) RAGE expression is up-regulated in human cerebral ischemia and pMCAO rats. *Neurosci. Lett.* **445**, 117–121
- 38 Menini, T., Ikeda, H., Kimura, S. and Gugliucci, A. (2014) Circulating soluble RAGE increase after a cerebrovascular event. *Clin. Chem. Lab. Med.* **52**, 109–116
- 39 Carnevale, D., Mascio, G., D'andrea, I., Fardella, V., Bell, R.D., Branchi, I. et al. (2012) Hypertension induces brain  $\beta$ -amyloid accumulation, cognitive impairment, and memory deterioration through activation of receptor for advanced glycation end products in brain vasculature. *Hypertension* **60**, 188–197

- 40 Geroldi, D., Falcone, C., Emanuele, E., D'Angelo, A., Calcagnino, M., Buzzi, M.P. et al. (2005) Decreased plasma levels of soluble receptor for advanced glycation end-products in patients with essential hypertension. *J. Hypertens.* **23**, 1725–1729
- 41 Haraba, R., Suica, V.I., Uyy, E., Ivan, L. and Antohe, F. (2011) Hyperlipidemia stimulates the extracellular release of the nuclear high mobility group box 1 protein. *Cell Tissue Res.* **346**, 361–368
- 42 Volz, H.C., Seidel, C., Laohachewin, D., Kaya, Z., Müller, O.J., Pleger, S.T. et al. (2010) HMGB1: the missing link between diabetes mellitus and heart failure. *Basic Res. Cardiol.* **105**, 805–820
- 43 Yao, D. and Brownlee, M. (2010) Hyperglycemia-induced reactive oxygen species increase expression of the receptor for advanced glycation end products (RAGE) and RAGE ligands. *Diabetes* **59**, 249–255
- 44 Bucala, R., Makita, Z., Vega, G., Grundy, S., Koschinsky, T., Cerami, A. et al. (1994) Modification of low density lipoprotein by advanced glycation end products contributes to the dyslipidemia of diabetes and renal insufficiency. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 9441–9445
- 45 Hu, P., Lai, D., Lu, P., Gao, J. and He, H. (2012) ERK and Akt signaling pathways are involved in advanced glycation end product-induced autophagy in rat vascular smooth muscle cells. *Int. J. Mol. Med.* **29**, 613–618
- 46 Aliprantis, A.O., Yang, R.-B., Mark, M.R., Suggett, S., Devaux, B., Radolf, J.D. et al. (1999) Cell activation and apoptosis by bacterial lipoproteins through toll-like receptor-2. *Science* **285**, 736–739
- 47 Said-Sadier, N. and Ojcius, D.M. (2012) Alarmins, inflammasomes and immunity. *Biomed. J.* **35**, 437–449
- 48 Chang, Z. (2010) Important aspects of Toll-like receptors, ligands and their signaling pathways. *Inflamm. Res.* **59**, 791–808
- 49 Brasier, A.R. (2006) The NF-kappaB regulatory network. *Cardiovasc. Toxicol.* **6**, 111–130
- 50 Gesuete, R., Kohama, S.G. and Stenzel-Poore, M.P. (2014) Toll-like receptors and ischemic brain injury. *J. Neuropathol. Exp. Neurol.* **73**, 378–386
- 51 Brea, D., Blanco, M., Ramos-Cabrer, P., Moldes, O., Arias, S., Perez-Mato, M. et al. (2011) Toll-like receptors 2 and 4 in ischemic stroke: outcome and therapeutic values. *J. Cereb. Blood Flow Metab.* **31**, 1424–1431
- 52 Caso, J.R., Pradillo, J.M., Hurtado, O., Lorenzo, P., Moro, M.A. and Lizasoain, I. (2007) Toll-like receptor 4 is involved in brain damage and inflammation after experimental stroke. *Circulation* **115**, 1599–1608
- 53 Tang, S.C., Arumugam, T.V., Xu, X., Cheng, A., Mughal, M.R., Jo, D.G. et al. (2007) Pivotal role for neuronal Toll-like receptors in ischemic brain injury and functional deficits. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 13798–13803
- 54 Balosso, S., Liu, J., Bianchi, M.E. and Vezzani, A. (2014) Disulfide-containing high mobility group box-1 promotes N-methyl-D-aspartate receptor function and excitotoxicity by activating Toll-like receptor 4-dependent signaling in hippocampal neurons. *Antioxid. Redox Signal.* **21**, 1726–1740
- 55 Lehnardt, S., Massillon, L., Follett, P., Jensen, F.E., Ratan, R., Rosenberg, P.A. et al. (2003) Activation of innate immunity in the CNS triggers neurodegeneration through a Toll-like receptor 4-dependent pathway. *Proc. Natl. Acad. Sci. U.S.A.* **100**, 8514–8519
- 56 Rosenberg, G.A. (2002) Matrix metalloproteinases in neuroinflammation. *Glia* **39**, 279–291
- 57 Yenari, M.A., Kauppinen, T.M. and Swanson, R.A. (2010) Microglial activation in stroke: therapeutic targets. *Neurotherapeutics* **7**, 378–391
- 58 Candelario-Jalil, E., Yang, Y. and Rosenberg, G.A. (2009) Diverse roles of matrix metalloproteinases and tissue inhibitors of metalloproteinases in neuroinflammation and cerebral ischemia. *Neuroscience* **158**, 983–994
- 59 Rosenberg, G.A., Cunningham, L.A., Wallace, J., Alexander, S., Estrada, E.Y., Grossetete, M. et al. (2001) Immunohistochemistry of matrix metalloproteinases in reperfusion injury to rat brain: activation of MMP-9 linked to stromelysin-1 and microglia in cell cultures. *Brain Res.* **893**, 104–112
- 60 del Zoppo, G.J., Milner, R., Mabuchi, T., Hung, S., Wang, X., Berg, G.I. et al. (2007) Microglial activation and matrix protease generation during focal cerebral ischemia. *Stroke* **38**, 646–651
- 61 Pfefferkorn, T. and Rosenberg, G.A. (2003) Closure of the blood-brain barrier by matrix metalloproteinase inhibition reduces rtPA-mediated mortality in cerebral ischemia with delayed reperfusion. *Stroke* **34**, 2025–2030
- 62 Asahi, M., Asahi, K., Jung, J.C., del Zoppo, G.J., Fini, M.E. and Lo, E.H. (2000) Role for matrix metalloproteinase 9 after focal cerebral ischemia: effects of gene knockout and enzyme inhibition with BB-94. *J. Cereb. Blood Flow Metab.* **20**, 1681–1689
- 63 Walker, E.J. and Rosenberg, G.A. (2009) TIMP-3 and MMP-3 contribute to delayed inflammation and hippocampal neuronal death following global ischemia. *Exp. Neurol.* **216**, 122–131
- 64 Zhao, B.Q., Wang, S., Kim, H.Y., Storrie, H., Rosen, B.R., Mooney, D.J. et al. (2006) Role of matrix metalloproteinases in delayed cortical responses after stroke. *Nat. Med.* **12**, 441–445
- 65 Sapojnikova, N., Kartvelishvili, T., Asatiani, N., Zinkevich, V., Kalandadze, I., Gugutsidze, D. et al. (2014) Correlation between MMP-9 and extracellular cytokine HMGB1 in prediction of human ischemic stroke outcome. *Biochim. Biophys. Acta* **1842**, 1379–1384
- 66 Qiu, J., Xu, J., Zheng, Y., Wei, Y., Zhu, X., Lo, E.H. et al. (2010) High-mobility group box 1 promotes metalloproteinase-9 upregulation through Toll-like receptor 4 after cerebral ischemia. *Stroke* **41**, 2077–2082
- 67 Liu, W., Hendren, J., Qin, X.J., Shen, J. and Liu, K.J. (2009) Normobaric hyperoxia attenuates early blood-brain barrier disruption by inhibiting MMP-9-mediated occludin degradation in focal cerebral ischemia. *J. Neurochem.* **108**, 811–820
- 68 Liu, J., Jin, X., Liu, K.J. and Liu, W. (2012) Matrix metalloproteinase-2-mediated occludin degradation and caveolin-1-mediated claudin-5 redistribution contribute to blood-brain barrier damage in early ischemic stroke stage. *J. Neurosci.* **32**, 3044–3057
- 69 Yang, Y., Estrada, E.Y., Thompson, J.F., Liu, W. and Rosenberg, G.A. (2007) Matrix metalloproteinase-mediated disruption of tight junction proteins in cerebral vessels is reversed by synthetic matrix metalloproteinase inhibitor in focal ischemia in rat. *J. Cereb. Blood Flow Metab.* **27**, 697–709
- 70 Aoki, T., Sumii, T., Mori, T., Wang, X. and Lo, E.H. (2002) Blood-brain barrier disruption and matrix metalloproteinase-9 expression during reperfusion injury: mechanical versus embolic focal ischemia in spontaneously hypertensive rats. *Stroke* **33**, 2711–2717

- 71 Sumii, T. and Lo, E.H. (2002) Involvement of matrix metalloproteinase in thrombolysis-associated hemorrhagic transformation after embolic focal ischemia in rats. *Stroke* **33**, 831–836
- 72 Kaur, J., Zhao, Z., Klein, G.M., Lo, E.H. and Buchan, A.M. (2004) The neurotoxicity of tissue plasminogen activator? *J. Cereb. Blood Flow Metab.* **24**, 945–963
- 73 Zhao, B.-Q., Ikeda, Y., Ihara, H., Urano, T., Fan, W., Mikawa, S. et al. (2004) Essential role of endogenous tissue plasminogen activator through matrix metalloproteinase 9 induction and expression on heparin-produced cerebral hemorrhage after cerebral ischemia in mice. *Blood* **103**, 2610–2616
- 74 Dang, A.B., Tay, B.K., Kim, H.T., Nauth, A., Alfonso-Jaume, M.A. and Lovett, D.H. (2008) Inhibition of MMP2/MMP9 after spinal cord trauma reduces apoptosis. *Spine (Phila Pa 1976)* **33**, E576–9
- 75 Kalinina, N., Agrotis, A., Antropova, Y., DiVitto, G., Kanellakis, P., Kostolias, G. et al. (2004) Increased expression of the DNA-binding cytokine HMGB1 in human atherosclerotic lesions: role of activated macrophages and cytokines. *Arterioscler. Thromb. Vasc. Biol.* **24**, 2320–2325
- 76 Porto, A., Palumbo, R., Pieroni, M., Aprigliano, G., Chiesa, R., Sanvito, F. et al. (2006) Smooth muscle cells in human atherosclerotic plaques secrete and proliferate in response to high mobility group box 1 protein. *FASEB J.* **20**, 2565–2566
- 77 Rouhiainen, A., Imai, S., Rauvala, H. and Parkkinen, J. (2000) Occurrence of amphoterin (HMG1) as an endogenous protein of human platelets that is exported to the cell surface upon platelet activation. *Thromb. Haemost.* **84**, 1087–1094
- 78 Fiuza, C., Bustin, M., Talwar, S., Tropea, M., Gerstenberger, E., Shelhamer, J.H. et al. (2003) Inflammation-promoting activity of HMGB1 on human microvascular endothelial cells. *Blood* **101**, 2652–2660
- 79 Treutiger, C.J., Mullins, G.E., Johansson, A.S., Rouhiainen, A., Rauvala, H.M., Erlandsson-Harris, H. et al. (2003) High mobility group 1 B-box mediates activation of human endothelium. *J. Intern. Med.* **254**, 375–385
- 80 Richard, S.A., Min, W., Su, Z. and Xu, H.-X. (2017) Epochal neuroinflammatory role of high mobility group box 1 in central nervous system diseases. *AIMS Mol. Sci.* **4**, 185–218
- 81 Inoue, K., Kawahara, K., Biswas, K.K., Ando, K., Mitsudo, K., Nobuyoshi, M. et al. (2007) HMGB1 expression by activated vascular smooth muscle cells in advanced human atherosclerosis plaques. *Cardiovasc. Pathol.* **16**, 136–143
- 82 Kanellakis, P., Agrotis, A., Kyaw, T.S., Koulis, C., Ahrens, I., Mori, S. et al. (2011) High-mobility group box protein 1 neutralization reduces development of diet-induced atherosclerosis in apolipoprotein e-deficient mice. *Arterioscler. Thromb. Vasc. Biol.* **31**, 313–319
- 83 Ito, T., Kawahara, K., Nakamura, T., Yamada, S., Nakamura, T., Abeyama, K. et al. (2007) High-mobility group box 1 protein promotes development of microvascular thrombosis in rats. *J. Thromb. Haemost.* **5**, 109–116
- 84 Richard, S.A., Min, W., Su, Z. and Xu, H. (2017) High mobility group box 1 and traumatic brain injury. *J. Behav. Brain Sci.* **7**, 50
- 85 Zhang, Z.G., Zhang, L., Jiang, Q., Zhang, R., Davies, K., Powers, C. et al. (2000) VEGF enhances angiogenesis and promotes blood-brain barrier leakage in the ischemic brain. *J. Clin. Invest.* **106**, 829–838
- 86 Sun, Y., Jin, K., Xie, L., Childs, J., Mao, X.O., Logvinova, A. et al. (2003) VEGF-induced neuroprotection, neurogenesis, and angiogenesis after focal cerebral ischemia. *J. Clin. Invest.* **111**, 1843–1851
- 87 Schlueter, C., Weber, H., Meyer, B., Rogalla, P., Röser, K., Hauke, S. et al. (2005) Angiogenetic signaling through hypoxia: HMGB1: an angiogenetic switch molecule. *Am. J. Pathol.* **166**, 1259–1263
- 88 Hayakawa, K., Pham, L.D., Katusic, Z.S., Arai, K. and Lo, E.H. (2012) Astrocytic high-mobility group box 1 promotes endothelial progenitor cell-mediated neurovascular remodeling during stroke recovery. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 7505–7510
- 89 Gleichman, A.J. and Carmichael, S.T. (2014) Astrocytic therapies for neuronal repair in stroke. *Neurosci. Lett.* **565**, 47–52
- 90 Passalacqua, M., Patrone, M., Picotti, G., Del Rio, M., Sparatore, B., Melloni, E. et al. (1997) Stimulated astrocytes release high-mobility group 1 protein, an inducer of LAN-5 neuroblastoma cell differentiation. *Neuroscience* **82**, 1021–1028
- 91 Huttunen, H.J., Kuja-Panula, J., Sorci, G., Agneletti, A.L., Donato, R. and Rauvala, H. (2000) Coregulation of neurite outgrowth and cell survival by amphoterin and S100 proteins through receptor for advanced glycation end products (RAGE) activation. *J. Biol. Chem.* **275**, 40096–40105
- 92 Huttunen, H.J., Kuja-Panula, J. and Rauvala, H. (2002) Receptor for advanced glycation end products (RAGE) signaling induces CREB-dependent chromogranin expression during neuronal differentiation. *J. Biol. Chem.* **277**, 38635–38646
- 93 Aghayev, K., Bal, E., Rahimli, T., Mut, M., Balc, S., Vrionis, F. et al. (2012) Aquaporin-4 expression is not elevated in mild hydrocephalus. *Acta Neurochir. (Wien)* **154**, 753–759
- 94 Xiong, X.X., Gu, L.J., Shen, J., Kang, X.H., Zheng, Y.Y., Yue, S.B. et al. (2014) Probenecid protects against transient focal cerebral ischemic injury by inhibiting HMGB1 release and attenuating AQP4 expression in mice. *Neurochem. Res.* **39**, 216–224
- 95 Xiong, X., White, R.E., Xu, L., Yang, L., Sun, X., Zou, B. et al. (2013) Mitigation of murine focal cerebral ischemia by the hypocretin/orexin system is associated with reduced inflammation. *Stroke* **44**, 764–770
- 96 Iacovetta, C., Rudloff, E. and Kirby, R. (2012) The role of aquaporin 4 in the brain. *Vet. Clin. Pathol.* **41**, 32–44
- 97 Manley, G.T., Fujimura, M., Ma, T., Noshita, N., Filiz, F., Bollen, A.W. et al. (2000) Aquaporin-4 deletion in mice reduces brain edema after acute water intoxication and ischemic stroke. *Nat. Med.* **6**, 159–163
- 98 Hiroaki, Y., Tani, K., Kamegawa, A., Gyobu, N., Nishikawa, K., Suzuki, H. et al. (2006) Implications of the aquaporin-4 structure on array formation and cell adhesion. *J. Mol. Biol.* **355**, 628–639
- 99 Saadoun, S., Papadopoulos, M.C., Watanabe, H., Yan, D., Manley, G.T. and Verkman, A.S. (2005) Involvement of aquaporin-4 in astroglial cell migration and glial scar formation. *J. Cell Sci.* **118**, 5691–5698
- 100 Fukuda, A.M. and Badaut, J. (2012) Aquaporin 4: a player in cerebral edema and neuroinflammation. *J. Neuroinflammation* **9**, 279
- 101 Ekdahl, C.T., Kokaia, Z. and Lindvall, O. (2009) Brain inflammation and adult neurogenesis: the dual role of microglia. *Neuroscience* **158**, 1021–1029
- 102 Kim, I.D., Shin, J.H., Kim, S.W., Choi, S., Ahn, J., Han, P.L. et al. (2012) Intranasal delivery of HMGB1 siRNA confers target gene knockdown and robust neuroprotection in the postischemic brain. *Mol. Ther.* **20**, 829–839



- 103 Bederson, J.B., Pitts, L.H., Tsuji, M., Nishimura, M.C., Davis, R.L. and Bartkowski, H. (1986) Rat middle cerebral artery occlusion: evaluation of the model and development of a neurologic examination. *Stroke* **17**, 472–476
- 104 Huang, J., Liu, B., Yang, C., Chen, H., Eunice, D. and Yuan, Z. (2013) Acute hyperglycemia worsens ischemic stroke-induced brain damage via high mobility group box-1 in rats. *Brain Res.* **1535**, 148–155
- 105 Li, P.A., Shuaib, A., Miyashita, H., He, Q.P., Siesjo, B.K. and Warner, D.S. (2000) Hyperglycemia enhances extracellular glutamate accumulation in rats subjected to forebrain ischemia. *Stroke* **31**, 183–192
- 106 Ennis, S.R. and Keep, R.F. (2007) Effect of sustained-mild and transient-severe hyperglycemia on ischemia-induced blood-brain barrier opening. *J. Cereb. Blood Flow Metab.* **27**, 1573–1582
- 107 Kamada, H., Yu, F., Nito, C. and Chan, P.H. (2007) Influence of hyperglycemia on oxidative stress and matrix metalloproteinase-9 activation after focal cerebral ischemia/reperfusion in rats: relation to blood-brain barrier dysfunction. *Stroke* **38**, 1044–1049
- 108 Cipolla, M.J., Huang, Q. and Sweet, J.G. (2011) Inhibition of protein kinase C $\beta$  reverses increased blood-brain barrier permeability during hyperglycemic stroke and prevents edema formation in vivo. *Stroke* **42**, 3252–3257
- 109 Shichita, T., Sugiyama, Y., Ooboshi, H., Sugimori, H., Nakagawa, R., Takada, I. et al. (2009) Pivotal role of cerebral interleukin-17-producing gammadeltaT cells in the delayed phase of ischemic brain injury. *Nat. Med.* **15**, 946–950
- 110 Liesz, A., Zhou, W., Mracsko, E., Karcher, S., Bauer, H., Schwarting, S. et al. (2011) Inhibition of lymphocyte trafficking shields the brain against deleterious neuroinflammation after stroke. *Brain* **134**, 704–720
- 111 Bodhankar, S., Chen, Y., Vandenbark, A.A., Murphy, S.J. and Offner, H. (2014) Treatment of experimental stroke with IL-10-producing B-cells reduces infarct size and peripheral and CNS inflammation in wild-type B-cell-sufficient mice. *Metab. Brain Dis.* **29**, 59–73
- 112 Liesz, A., Suri-Payer, E., Veltkamp, C., Doerr, H., Sommer, C., Rivest, S. et al. (2009) Regulatory T cells are key cerebroprotective immunomodulators in acute experimental stroke. *Nat. Med.* **15**, 192–199
- 113 Ren, X., Akiyoshi, K., Dziennis, S., Vandenbark, A.A., Herson, P.S., Hurn, P.D. et al. (2011) Regulatory B cells limit CNS inflammation and neurologic deficits in murine experimental stroke. *J. Neurosci.* **31**, 8556–8563
- 114 Bodhankar, S., Chen, Y., Vandenbark, A.A., Murphy, S.J. and Offner, H. (2013) IL-10-producing B-cells limit CNS inflammation and infarct volume in experimental stroke. *Metab. Brain Dis.* **28**, 375–386
- 115 Liesz, A., Zhou, W., Na, S.Y., Hammerling, G.J., Garbi, N., Karcher, S. et al. (2013) Boosting regulatory T cells limits neuroinflammation in permanent cortical stroke. *J. Neurosci.* **33**, 17350–17362
- 116 Ooboshi, H., Ibayashi, S., Shichita, T., Kumai, Y., Takada, J., Ago, T. et al. (2005) Postischemic gene transfer of interleukin-10 protects against both focal and global brain ischemia. *Circulation* **111**, 913–919
- 117 Whiteley, W., Jackson, C., Lewis, S., Lowe, G., Rumley, A., Sandercock, P. et al. (2009) Inflammatory markers and poor outcome after stroke: a prospective cohort study and systematic review of interleukin-6. *PLoS Med.* **6**, e1000145
- 118 Andersson, U., Wang, H., Palmblad, K., Aveberger, A.C., Bloom, O., Erlandsson-Harris, H. et al. (2000) High mobility group 1 protein (HMG-1) stimulates proinflammatory cytokine synthesis in human monocytes. *J. Exp. Med.* **192**, 565–570
- 119 Ren, D., Sun, R. and Wang, S. (2006) Role of inducible nitric oxide synthase expressed by alveolar macrophages in high mobility group box 1–induced acute lung injury. *Inflamm. Res.* **55**, 207–215
- 120 Iadecola, C., Zhang, F. and Xu, X. (1995) Inhibition of inducible nitric oxide synthase ameliorates cerebral ischemic damage. *Am. J. Physiol.* **268**, R286–92
- 121 Barome, F., Aravin, B. and White, R. (1997) Tumor necrosis factor- $\alpha$ : a mediator of focal ischemic brain injury. *Stroke* **28**, 1233–1244
- 122 Gong, G., Xiang, L., Yuan, L., Hu, L., Wu, W., Cai, L. et al. (2014) Protective effect of glycyrrhizin, a direct HMGB1 inhibitor, on focal cerebral ischemia/reperfusion-induced inflammation, oxidative stress, and apoptosis in rats. *PLoS ONE* **9**, e89450
- 123 Lo, E.H., Dalkara, T. and Moskowitz, M.A. (2003) Mechanisms, challenges and opportunities in stroke. *Nat. Rev. Neurosci.* **4**, 399–415
- 124 Yang, J., Huang, C., Yang, J., Jiang, H. and Ding, J. (2010) Statins attenuate high mobility group box-1 protein induced vascular endothelial activation: a key role for TLR4/NF- $\kappa$ B signaling pathway. *Mol. Cell. Biochem.* **345**, 189–195
- 125 Wang, L., Zhang, X., Liu, L., Yang, R., Cui, L. and Li, M. (2010) Atorvastatin protects rat brains against permanent focal ischemia and downregulates HMGB1, HMGB1 receptors (RAGE and TLR4), NF- $\kappa$ B expression. *Neurosci. Lett.* **471**, 152–156
- 126 Parkkinen, J., Raulo, E., Merenmies, J., Nolo, R., Kajander, E.O., Baumann, M. et al. (1993) Amphoterin, the 30-kDa protein in a family of HMG1-type polypeptides. Enhanced expression in transformed cells, leading edge localization, and interactions with plasminogen activation. *J. Biol. Chem.* **268**, 19726–19738
- 127 Papadopoulos, M.C. and Verkman, A.S. (2008) Potential utility of aquaporin modulators for therapy of brain disorders. *Prog. Brain Res.* **170**, 589–601
- 128 Suadicani, S.O., Brosnan, C.F. and Scemes, E. (2006) P2X7 receptors mediate ATP release and amplification of astrocytic intercellular Ca $^{2+}$  signaling. *J. Neurosci.* **26**, 1378–1385
- 129 Silverman, W., Locovei, S. and Dahl, G. (2008) Probenecid, a gout remedy, inhibits pannexin 1 channels. *Am. J. Physiol. Cell Physiol.* **295**, C761–7
- 130 Dahl, G. and Keane, R.W. (2012) Pannexin: from discovery to bedside in 11+/-4 years? *Brain Res.* **1487**, 150–159
- 131 Hayakawa, K., Mishima, K., Irie, K., Hazekawa, M., Mishima, S., Fujioka, M. et al. (2008) Cannabidiol prevents a post-ischemic injury progressively induced by cerebral ischemia via a high-mobility group box1-inhibiting mechanism. *Neuropharmacology* **55**, 1280–1286
- 132 Hayakawa, K., Mishima, K., Nozaki, M., Hazekawa, M., Mishima, S., Fujioka, M. et al. (2008) Delayed treatment with minocycline ameliorates neurologic impairment through activated microglia expressing a high-mobility group box1-inhibiting mechanism. *Stroke* **39**, 951–958
- 133 Kikuchi, K., Kawahara, K., Tancharoen, S., Matsuda, F., Morimoto, Y., Ito, T. et al. (2009) The free radical scavenger edaravone rescues rats from cerebral infarction by attenuating the release of high-mobility group box-1 in neuronal cells. *J. Pharmacol. Exp. Ther.* **329**, 865–874
- 134 Ye, X., Chopp, M., Liu, X., Zacharek, A., Cui, X., Yan, T. et al. (2011) Niaspan reduces high-mobility group box 1/receptor for advanced glycation endproducts after stroke in type-1 diabetic rats. *Neuroscience* **190**, 339–345

- 135 Jiang, W.L., Xu, Y., Zhang, S.P., Zhu, H.B. and Hou, J. (2012) Tricin 7-glucoside protects against experimental cerebral ischemia by reduction of NF-kappaB and HMGB1 expression. *Eur. J. Pharm. Sci.* **45**, 50–57
- 136 Wang, L., Zhang, X., Liu, L., Cui, L., Yang, R., Li, M. et al. (2010) Tanshinone II A down-regulates HMGB1, RAGE, TLR4, NF-kappaB expression, ameliorates BBB permeability and endothelial cell function, and protects rat brains against focal ischemia. *Brain Res.* **1321**, 143–151
- 137 Gwak, G.Y., Moon, T.G., Lee, D.H. and Yoo, B.C. (2012) Glycyrrhizin attenuates HMGB1-induced hepatocyte apoptosis by inhibiting the p38-dependent mitochondrial pathway. *World J. Gastroenterol.* **18**, 679–684