

■ C O R R E C T I O N

The UKPDS risk engine: a model for the risk of coronary heart disease in Type II diabetes (UKPDS 56)

Richard J. STEVENS, Viti KOTHARI, Amanda I. ADLER, Irene M. STRATTON and Rury R. HOLMAN on behalf of the United Kingdom Prospective Diabetes Study (UKPDS) Group

Volume 101 (2001), pp. 671–679

The Publisher regrets that the following errors appeared in the above published paper:

Page 673, right column, line 11: for ‘where d is the duration of diagnosed diabetes’ *read* ‘where d is the effect parameter for the duration of diagnosed diabetes’.

Page 673, right column, line 18: for ‘The survival probability for t years in a patient who has had diabetes for T years is also useful’ *read* ‘The risk over t years in a patient who has had diabetes for T years is also useful’.

Page 673, right column, line 23: for $(t-1, t)$ *read* $(t-1, t]$.

Page 675, Figure 2 legend: for ‘(a) HbA_{1c} value’ *read* ‘(a) By HbA_{1c} value’, for ‘(b) Systolic blood pressure’ *read* ‘(b) By systolic blood pressure’, and for ‘(c) Lipid ratio’ *read* ‘(c) By lipid ratio’.

Page 675, left column, lines 6 and 8: for 0.112 *read* 0.0112.

Page 675, left column, line 12: for 0.0883 *read* 0.00883.

Page 675, right column, line 33: for 4541 *read* 4540.

Page 677, right column, line 27: for 1.22 *read* 1.23.

Page 677, right column, line 31: for $3.845^{1.22} = 5.171$ *read* $3.845^{1.23} = 5.24$.

Page 678, left column, line 20: for ‘ $P(t) = 0.049$ when qd^{t-1} ’ *read* ‘ $P(t) = 0.049$ when $qd^{t-1} = 0.05$ ’.

Page 678, right column, reference 8: for ‘Scandinavian Simvastatin (4S)’ *read* ‘Scandinavian Simvastatin Survival Study (4S)’.

Page 679, left column, reference 28: for <http://hyp.ac.uk/bhs/managmt.html>

read <http://www.hyp.ac.uk/bhs/managemt.html>